

FORM PTO-1390
(REV. 11-2000)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

SF0820K2

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

09787192

INTERNATIONAL APPLICATION NO.

INTERNATIONAL FILING DATE

PRIORITY DATE CLAIMED

PCT/US99/22269

23 September 1999

25 September 1998

TITLE OF INVENTION

ANTIBODIES TO MAMMALIAN LANGERHANS CELL ANTIGEN AND THEIR USES

APPLICANT(S) FOR DO/EO/US DUVERT-FRANCES, Valerie, PIN, Jean-Jacques, VALLADEAU, Jenny, CLAIR, Valerie, SAELAND, Sem and LEBECQUE, Serge J. E.

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below.
4. ☒ The US has been elected by the expiration of 19 months from the priority date (Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ has been communicated by the International Bureau.
 - c. ☒ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
 - a. ☐ is attached hereto.
 - b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ have been communicated by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11 to 20 below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A **FIRST** preliminary amendment.
14. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
15. ☐ A substitute specification.
16. ☐ A change of power of attorney and/or address letter.
17. ☐ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.
18. ☐ A second copy of the published international application under 35 U.S.C. 154(d)(4).
19. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
20. ☐ Other items or information:

21. ☐ The following fees are submitted:**BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)):**

Neither international preliminary examination fee (37 CFR 1.482)
nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO
and International Search Report not prepared by the EPO or JPO. \$1000.00

International preliminary examination fee (37 CFR 1.482) not paid to
USPTO but International Search Report prepared by the EPO or JPO. \$860.00

International preliminary examination fee (37 CFR 1.482) not paid to USPTO
but international search fee (37 CFR 1.445(a)(2)) paid to USPTO. \$710.00

International preliminary examination fee (37 CFR 1.482) paid to USPTO
but all claims did not satisfy provisions of PCT Article 33(1)-(4). \$690.00

International preliminary examination fee (37 CFR 1.482) paid to USPTO
and all claims satisfied provisions of PCT Article 33(1)-(4). \$100.00

ENTER APPROPRIATE BASIC FEE AMOUNT =**CALCULATIONS PTO USE ONLY**

Surcharge of \$130.00 for furnishing the oath or declaration later than ☐ 20 ☐ 30
months from the earliest claimed priority date (37 CFR 1.492(e)).

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	15 - 20 =		x \$18.00

Independent claims	11 - 3 =	8	x \$80.00
--------------------	----------	---	-----------

MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$270.00
---	--	--	------------

TOTAL OF ABOVE CALCULATIONS =

☐ Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above
are reduced by 1/2.

SUBTOTAL =

Processing fee of \$130.00 for furnishing the English translation later than ☐ 20 ☐ 30
months from the earliest claimed priority date (37 CFR 1.492(f)).

TOTAL NATIONAL FEE =

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be
accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +

TOTAL FEES ENCLOSED =

Amount to be refunded:	\$
---------------------------	----

charged:	\$
----------	----

- a. ☐ A check in the amount of \$ _____ to cover the above fees is enclosed.
- b. ☒ Please charge my Deposit Account No. 9-0365 _____ in the amount of \$ 1,500.00 to cover the above fees.
A duplicate copy of this sheet is enclosed.
- c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any
overpayment to Deposit Account No. 9-0365 _____. A duplicate copy of this sheet is enclosed.
- d. ☐ Fees are to be charged to a credit card. **WARNING:** Information on this form may become public. **Credit card
information should not be included on this form.** Provide credit card information and authorization on PTO-2038.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137 (a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

Jaye P. McLaughlin
Patent Dept. K-6-1, 1990

Schering-Plough Corporation

2000 Galloping Hill Road

Kenilworth, New Jersey 07033-0530



SIGNATURE

Jaye P. McLaughlin

NAME

41211

REGISTRATION NUMBER

Express Mail Label no. EL 664530682 US

SF0820K2

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

-----X
 In re Application of:
 Duvert-Frances et al.

:
 : Examiner: To Be Assigned
 : Group Art Unit: To Be Assigned

For Patent For:

ANTIBODIES TO MAMMALIAN
 LANGERHANS CELL ANTIGEN AND
 THEIR USES

:

Serial No.:

: Date: March 15, 2001

Filed:

:

-----X
 Schering-Plough Corporation
 Kenilworth, New Jersey 07033-0530

Honorable Commissioner of Patents
 and Trademarks
 Washington, D.C. 20231

Sir:

PRELIMINARY AMENDMENT

Please amend the above-identified application as follows.

IN THE SPECIFICATION

On page 1 of the specification, underneath the title, please enter the following statement:

--The present application is a 35 U.S.C. 371 National Phase application of PCT/US99/22269, filed September 23, 1999, which claims priority from European Application Number 98 402 374.7, filed September 25, 1998, and European Application Number 99 400 394.5, filed February 18, 1999.--

EXPRESS MAIL NO. EL 664530682 US

09787192-034501

SCHERING-PLOUGH CORPORATION

Patent Department, K-6-1, 1990

2000 Galloping Hill Road

Kenilworth, NJ 07033-0530

908-298-4299

Date: March 15, 2001

Respectfully submitted,



Jaye R. McLaughlin

Reg. No. 41,211

Attorney for Applicants

(908) 298-4299

10586_1

09787492.031504
FOSTER 26128260

ANTIBODIES TO MAMMALIAN DC ANTIGEN AND USES

5 FIELD OF THE INVENTION

The present invention relates to compositions which function in controlling physiology, development, and differentiation of mammalian cells, e.g., cells of a mammalian immune system. In particular, it provides 10 antibodies, e.g., agonists and antagonists, which regulate cellular physiology, development, differentiation, or function of various cell types, including hematopoietic cells, and particularly dendritic cells e.g., Langerhans cells.

15 BACKGROUND OF THE INVENTION

Dendritic cells (DC) are antigen-presenting cells that are required for initiation of a specific immune response. See, e.g., Banchereau and Steinman (1998) 20 Nature 392:242-52. One type of DC is exemplified by Langerhans cells (LC), immature DC cells that reside in non-lymphoid tissue, such as the epidermis, and whose primary function is to capture antigen. See, e.g., Steinman, et al. (1995) J. Exp. Med. 182:283-288.

25 Antigen capture is achieved primarily through specialized surface-membrane endocytic structures or through macropinocytosis, thus permitting Langerhans cells to concentrate solutes which are present in large volumes of fluid. See, e.g., Sallusto, et al. (1995) J. Exp. Med. 182:389-400. Concomitant with processing of 30 antigen in specialized organelles of the endocytic pathway, DC, such as Langerhans cells, migrate to secondary lymphoid tissue and undergo a number of phenotypic modifications. These maturation events 35 ultimately translate into highly efficient presentation of processed antigen, by appropriate MHC molecules of the DC, to T cells.

In particular, the maturation process of the DC Langerhans cell includes loss of adhesion receptors such

09787192-031501

as E-cadherin, and the disappearance of Birbeck granules (BG), which are characteristic for LC. Conversely, upon acquisition of antigen-presentation function, costimulatory receptors such as the CD80 and CD86

- 5 molecules are upregulated on DC to permit T cell activation. Maturation events of DC can be reconstituted in vitro by TNF- α and CD40-ligand which mimic, respectively, the response to pro-inflammatory cytokines following encounter with pathogen, and the response to
10 contact with T cells in secondary lymphoid tissue. See, e.g., Caux, et al. (1996) J. Exp. Med. 184:695-706.

Thus, it is apparent that the highly specialized and anatomically localized functions of DC are controlled by tight regulation of the expression of a number of key
15 molecules.

- Recently, culture systems have become available to obtain large numbers of DC when cultured in the presence of cytokines. Using such culture methods, DC can be obtained for in vitro study from CD34⁺ hematopoietic
20 progenitor cells (HPC) present in cord blood when the HPC are co-cultured with TNF- α and GM-CSF (such cultures are referred to as CD34-derived DC), or from peripheral blood monocytes when they are co-cultured with GM-CSF and IL-4 (such cultures are referred to as monocyte-derived DC).
25 See, e.g., Caux, et al. (1992) Nature 392:258-261; Chapuis, et al. (1997) Eur. J. Immunol. 27:431-441; Romani, et al. (1994) J. Invest. Dermatol. 93:600-609.

- These methods permit either, the in vitro (see, e.g., Caux, et al. (1996) J. Exp. Med. 184:695-706), or,
30 the ex-vivo (from various organs; see e.g., Grouard, et al. (1996) Nature 384:364-367; O'Doherty, et al. (1994) Immunol. 82:487-493; Zhou (1995) J. Immunol. 154:3821-3835) isolation of phenotypically and functionally distinct DC subpopulations.

- 35 Consequently, it would be of great benefit to possess novel reagents capable of identifying markers that are expressed and associated with different DC subpopulations. These markers are detectable using antibodies, e.g., monoclonal or polyclonal. Such markers

09737162.031504

would permit the monitoring, characterization, and/or isolation of defined subsets of immature DCs by facilitating, e.g., cell-sorting and functional studies. Thus, needs exist for tools which permit a better understanding of the molecules involved in DC maturation, antigen presentation, and the mechanisms of DC interaction with other molecules, cells, and tissues. The present invention fulfills such needs by providing useful reagents and compositions involved in DC maturation and function.

SUMMARY OF THE INVENTION

The present invention is based, in part, upon the discovery of an antibody which defines and recognizes a novel cell antigen found on dendritic cells (DC) and Langerhans cells (LC). This monoclonal antibody is designated DCGM4 and the antigen it recognizes has been designated Langerin. The invention embraces this antibody and methods for its use. In addition, the invention is directed to antigen recognized by this antibody, along with variants of these proteins, e.g., mutations (muteins) of the natural sequence, species and allelic variants, fusion proteins, chemical mimetics, and other structural or functional analogs. Various uses of these different antibodies and protein compositions are also provided.

The present invention provides antibody which binds specifically to a mammalian Langerin. In preferred embodiments the mammal is a primate; or the antibodies are a monoclonal antibody, interfere with binding of DCGM4 to the Langerin, or are detectably labeled, e.g., with a fluorescent or enzymatic label.

The invention also provides methods of detecting a mammalian Langerin, comprising binding DCGM4 to Langerin. In various embodiments, the antibody is a labeled antibody or is immobilized to a solid substrate; the Langerin is expressed on a cell surface; the detecting allows isolation of a cell which comprises a nucleic acid which expresses Langerin; or the detecting further allows

00787192-031501

purification of Langerin. The invention also embraces a kit for detecting Langerin with a compartment containing an antibody. In preferred embodiments, the kit is a fluorescence immunoassay kit.

5 The present invention further provides methods of modulating an immune function modulated by a cell comprising contacting said cell with an antibody described herein. For instance, the modulation can be blocking DC cell maturation or function.

10 Also embraced herein are methods for analyzing a DC cell population, comprising measuring the presence of Langerin. Typically, the measuring is a quantitative determination, e.g., by measuring binding of an antibody to Langerin.

15 The invention also provides substantially pure mammalian Langerin antigens. Human Langerin and mouse Langerin are specifically described. Langerin can be purified by, for example, immunoaffinity, e.g., using an antibody which binds specifically to a mammalian
20 Langerin. A preferred antibody for such is DCGM4. Along with full length Langerin, the invention provides fragments which express an immunological epitope of said Langerin or modulate an immune response, e.g., a response mediated by a DC cell, including a Langerin⁺ cell.

25 The invention further provides genomic DNA of human langerin.

DETAILED DESCRIPTION OF THE INVENTION

30 All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety including all figures, graphs, and drawings.

35 General

 The present invention provides antibodies which recognize a mammalian protein that exhibits properties characteristic of functionally significant DC expressed

09707192.031504

molecules. The antibody is exemplified in one embodiment by a monoclonal antibody designated DCGM4.

The mammalian protein defined, e.g., selectively recognized, by the antibody DCGM4 is designated Langerin. The natural Langerin protein mediates various physiological responses leading to biological or physiological responses in target cells. In particular, Langerhans DC cells are responsible for antigen presentation, e.g., presentation of haptens in hypersensitivity reactions. The DCGM4 antibody modulates various immunological responses which affect DC maturation and antigen presentation.

Langerin is a 40 kDa N-glycosylated protein. Immunoprecipitation with DCGM4 from DC extracts and, subsequent elution with SDS-PAGE sample buffer yielded a homogeneous band of 40-42 kDa molecular mass. If DTT was omitted all along the purification steps, the profile was not modified on the gel, suggesting that Langerin is present at the cell membrane as a single chain or as an homodimer with non-covalent association. Two dimensional analysis confirmed the molecular mass of the molecule and indicated a pI of 5.2-5.5. Finally, Langerin is a glycoprotein, and most of the carbohydrate constituents were removed by N-glycosylase treatment.

A gene encoding human langerin has now been cloned. The nucleotide sequence is showing in SEQ ID NO:1. The predicted amino acid sequence is shown in SEQ ID NO: 2. The predicted protein is a type II membrane lectin, with a calcium-dependent carbohydrate recognition domain. Amino acid homology with the Kupffer receptor of mouse, rat and chicken was found and provides support that langerin functions as a langerhans cell specific receptor for antigen capture. The carbohydrate recognition domain motif EPN supports that the sugar moieties are probably mannose and glucose.

Identifying characteristics that permit one of ordinary skill in the art to distinguish Langerin from other proteins are listed below.

- binds with specificity to DCGM4 monoclonal antibody

- N-glycosylated form of protein ~40 kD, both reduced or non-reduced with a pI of 5.2-5.5 and lacking interchain disulfide bonds

5

- expressed on subset of DC cells
- normally expressed on the cell surface of Langerhans cells

10

- after binding with DCGM4, Langerin is transported within endocytic coated pits and participates in cytomembrane sandwiching; subsequently Langerin is associated with Birbeck granules.

15

Cells of the clonally derived hybridoma cell line designated DCGM4 from which mAb DCGM4 has been purified have been deposited with the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas VA, USA, on September 22, 1998 under ATCC Accession

20

Number No. HB-12576.

It is to be understood that this invention is not limited to the particular methods, compositions and antibodies described herein, as such methods, compositions and antibodies may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which is only limited by the appended claims.

30

II. Antibodies

Antibodies can be raised to the various mammalian, e.g., primate Langerin proteins and fragments thereof, both in naturally occurring native forms and in their recombinant forms, the difference being that antibodies to the active Langerin are more likely to recognize epitopes which are only present in the native conformations. Denatured antigen detection can also be

35

09787162-0115111

useful in, e.g., Western analysis. Anti-idiotypic antibodies are also contemplated, which would be useful as agonists or antagonists of a natural Langerin protein or an antibody.

- 5 Antibodies, including binding fragments and single chain versions, against predetermined fragments or the whole of the protein (e.g., a protein having an amino acid sequence shown in SEQ ID NO: 2) can be raised by immunization of animals. Monoclonal antibodies are
10 prepared from cells secreting the desired antibody. These antibodies can be screened for binding to normal or defective protein, or screened for agonistic or antagonistic activity. These monoclonal antibodies will usually bind with at least a K_D of about 1 mM, more
15 usually at least about 300 μ M, typically at least about 100 μ M, more typically at least about 30 μ M, preferably at least about 10 μ M, and more preferably at least about 3 μ M or better.

- 20 The antibodies, exemplified by the DCGM4, including antigen binding fragments, can have significant diagnostic or therapeutic value. They can be potent antagonists that bind to Langerin and inhibit binding partner interaction or inhibit the ability of the interaction to mediate a biological response. They also
25 can be useful as non-neutralizing antibodies and can be coupled to toxins or radionuclides so that when the antibody binds to the antigen, a cell expressing it, e.g., on its surface, is killed. Further, this antibody can be conjugated to drugs or other therapeutic agents,
30 either directly or indirectly by means of a linker, and may effect drug targeting.

- 35 The antibodies of this invention can also be useful in diagnostic applications. As capture or non-neutralizing antibodies, they might bind Langerin without inhibiting Langerin function on cells of the immune system, for example immature DC. As neutralizing antibodies, they can be useful in competitive binding assays. They will also be useful in detecting or quantifying Langerin protein or its binding partners.

0957192-034504

They may be used as reagents for Western blot analysis, or for immunoprecipitation or immunopurification of the Langerin protein. They will also be useful in evaluating cell populations to determine, e.g., the physiological state of an immune system.

Antigen fragments may be joined to other materials, particularly polypeptides, as fused or covalently joined polypeptides to be used as immunogens. The antigen may be purified as described below, including immunoaffinity methods using antibodies, e.g., DCGM4. An antigen and its fragments may be fused or covalently linked to a variety of carriers, such as keyhole limpet hemocyanin, bovine serum albumin, tetanus toxoid, etc. See, e.g., Microbiology, Hoeber Medical Division, Harper and Row, 1969; Landsteiner (1962) Specificity of Serological Reactions, Dover Publications, New York; and Williams, et al. (1967) Methods in Immunology and Immunochemistry, Vol. 1, Academic Press, New York, each of which are incorporated herein by reference, for descriptions of methods of preparing polyclonal antisera. A typical method involves hyperimmunization of an animal with the antigen. The blood of the animal is then collected shortly after the repeated immunizations and the gamma globulin is isolated.

In some instances, it is desirable to prepare monoclonal antibodies from various mammalian hosts, such as mice, rodents, primates, humans, etc. Description of techniques for preparing such monoclonal antibodies may be found in, e.g., Stites, et al. (eds.) Basic and Clinical Immunology (4th ed.), Lange Medical Publications, Los Altos, CA, and references cited therein; Harlow and Lane (1988) Antibodies: A Laboratory Manual, CSH Press; Goding (1986) Monoclonal Antibodies: Principles and Practice (2d ed.) Academic Press, New York; and particularly in Kohler and Milstein (1975) in Nature 256: 495-497, which discusses one method of generating monoclonal antibodies. Each of these references is incorporated herein by reference. Summarized briefly, this method involves injecting an

09787192-031504

animal with an immunogen. The animal is then sacrificed and cells taken from its spleen, which are then fused with myeloma cells. The result is a hybrid cell or "hybridoma" that is capable of reproducing in vitro. The population of hybridomas is then screened to isolate individual clones, each of which secrete a single antibody species to the immunogen. In this manner, the individual antibody species obtained are the products of immortalized and cloned single B cells from the immune animal generated in response to a specific site recognized on the immunogenic substance.

Other suitable techniques involve in vitro exposure of lymphocytes to the antigenic polypeptides or, alternatively, selection of libraries of antibodies in phage or similar vectors. See, Huse, et al. (1989) "Generation of a Large Combinatorial Library of the Immunoglobulin Repertoire in Phage Lambda," Science 246:1275-1281; and Ward, et al. (1989) Nature 341:544-546, each of which is hereby incorporated herein by reference. The polypeptides and antibodies of the present invention may be used with or without modification, including chimeric or humanized antibodies. Frequently, the polypeptides and antibodies will be labeled by joining, either covalently or non-covalently, a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, chemiluminescent moieties, magnetic particles, and the like. Patents teaching the use of such labels include U.S. Patent Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241. Also, recombinant immunoglobulins may be produced, see Cabilly, U.S. Patent No. 4,816,567; or made in transgenic mice, see Mendez, et al. (1997) Nature Genetics 15:146-156. These patents are incorporated herein by reference.

09787192.034501

The antibodies of this invention can also be used for affinity chromatography in isolating the protein. Columns can be prepared where the antibodies are linked to a solid support, e.g., particles, such as agarose, Sephadex, or the like, where a cell lysate may be passed through the column, the column washed, followed by increasing concentrations of a mild denaturant, whereby the purified Langerin protein will be released. Alternatively, the antibody may be used to quantitate and identify fractionated samples containing the antigen. Standard protein purification procedures, e.g., chromatography, will be used to enrich and purify Langerin protein with, e.g., ELISA assays, to identify fractions where Langerin separates.

Purified protein will be sequenced. The sequence will allow selection of oligonucleotide sequences useful as primers or probes. Alternatively, the sequence allows production of polypeptide segments for making additional antibodies. In addition, purified protein may be used for immunization, allowing production of polyclonal or monoclonal antibodies.

The antibodies may also be used to screen expression libraries for particular expression products. Usually the antibodies used in such a procedure will be labeled with a moiety allowing easy detection of presence of antigen by antibody binding. This will allow isolation of a cell which expresses a nucleic acid, e.g., a vector, encoding the antigen by, e.g., fluorescence activated cell sorting (FACS) analysis, and enrichment. Alternatively, an affinity method using antibodies of this invention can be used to immobilize and separate cells expressing the Langerin, e.g., encoded on a vector.

Antibodies raised against each Langerin protein will also be useful to raise anti-idiotypic antibodies. These will be useful in detecting or diagnosing various immunological conditions related to expression of the respective antigens.

III. Purified Langerin protein

Human Langerin protein can be isolated from natural sources using standard biochemical purification techniques and/or by use of the antibody to determine the presence of the antigen in particular fractionation

- 5 procedures. The purified proteins allow both sequence determination and preparation of peptides to generate further antibodies to recognize such segment. As used herein, Langerin shall encompass, when used in a protein context, a protein which, in a natural state, exhibits
- 10 the properties listed above, or a significant fragment of such a protein. e.g., a protein having an amino acid sequence set forth in SEQ ID NO: 2 or a subsequence thereof. It also refers to a mammalian, e.g., primate, derived polypeptide which exhibits similar biological
- 15 function or interacts with Langerin protein specific binding components. These binding components, e.g., antibodies, typically bind to a Langerin protein with high affinity, e.g., at least about 100 nM, usually better than about 30 nM, preferably better than about 10
- 20 nM, and more preferably at better than about 3 nM. One such preferred binding component is the antibody DCGM4.

The purified protein or peptide fragments are useful for generating antibodies by standard methods, as described below. Synthetic peptides or purified protein

25 can be presented to an immune system to generate a specific binding composition, e.g., monoclonal or polyclonal antibodies. See, e.g., Coligan (1991) Current Protocols in Immunology Wiley/Greene; and Harlow and Lane (1989) Antibodies: A Laboratory Manual Cold Spring Harbor

30 Press.

The term polypeptide, as used herein, includes a significant fragment or segment, and encompasses a stretch of amino acid residues of at least about 8 amino acids, generally at least 10 amino acids, more generally

35 at least 12 amino acids, often at least 14 amino acids, more often at least 16 amino acids, typically at least 18 amino acids, more typically at least 20 amino acids, usually at least 22 amino acids, more usually at least 24 amino acids, preferably at least 26 amino acids, more

preferably at least 28 amino acids, and, in particularly preferred embodiments, at least about 30 or more amino acids. Preferably, the fragment exhibits a biological property in common with the full length Langerin, e.g., immunological activity, including sharing of an epitope.

Substantially pure, in the polypeptide context, typically means that the protein is free from other contaminating proteins, nucleic acids, and other biologicals derived from the original source organism.

Purity may be assayed by standard methods, and will ordinarily be at least about 40% pure, more ordinarily at least about 50% pure, generally at least about 60% pure, more generally at least about 70% pure, often at least about 75% pure, more often at least about 80% pure, typically at least about 85% pure, more typically at least about 90% pure, preferably at least about 95% pure, more preferably at least about 98% pure, and in most preferred embodiments, at least 99% pure. The analysis may be weight or molar percentages, evaluated, e.g., by gel staining, spectrophotometry, or terminus labeling.

A binding composition or agent refers to molecules that bind with specificity to Langerin protein, e.g., in a ligand-receptor type fashion, an antibody-antigen interaction, or compounds, e.g., proteins which specifically associate with Langerin protein, e.g., in a natural physiologically relevant protein-protein interaction, either covalent or non-covalent. The molecule may be a polymer, or chemical reagent. This implies both binding affinity and binding specificity or selectivity. A functional analog may be a protein with structural modifications, or may be a wholly unrelated molecule, e.g., which has a molecular shape which interacts with the appropriate binding determinants. The proteins may serve as agonists or antagonists of a receptor, see, e.g., Goodman, et al. (eds. 1990) Goodman & Gilman's: The Pharmacological Bases of Therapeutics (8th ed.) Pergamon Press, Tarrytown, N.Y.

Soluble fragments of both the antibodies and Langerin antigens are provided by the invention.

09787192-031501

Solubility of a polypeptide or fragment depends upon the environment and the polypeptide. Many parameters affect polypeptide solubility, including temperature, electrolyte environment, size and molecular

- 5 characteristics of the polypeptide, and nature of the solvent. Typically, the temperature at which the polypeptide is used ranges from about 4° C to about 65° C. Usually the temperature at use is greater than about 18° C and more usually greater than about 22° C. For
10 diagnostic purposes, the temperature will usually be about room temperature or warmer, but less than the denaturation temperature of components in the assay. For therapeutic purposes, the temperature will usually be body temperature, typically about 37° C for humans,
15 though under certain situations the temperature may be raised or lowered in situ or in vitro.

The electrolytes will usually approximate in situ physiological conditions, but may be modified to higher or lower ionic strength where advantageous. The actual
20 ions may be modified, e.g., to conform to standard buffers used in physiological or analytical contexts.

The size and structure of the polypeptide should generally be in a substantially stable state, and usually not in a denatured state. The polypeptide may be
25 associated with other polypeptides in a quaternary structure, e.g., to confer solubility, or associated with lipids or detergents in a manner which approximates natural lipid bilayer interactions.

The solvent will usually be a biologically
30 compatible buffer, of a type used for preservation of biological activities, and will usually approximate a physiological solvent. Usually the solvent will have a neutral pH, typically between about 5 and 10, and preferably about 7.5. On some occasions, a detergent
35 will be added, typically a mild non-denaturing one, e.g., CHS or CHAPS, or a low enough concentration as to avoid significant disruption of structural or physiological properties of the antigen.

09787492.031501

Solubility is reflected by sedimentation measured in Svedberg units, which are a measure of the sedimentation velocity of a molecule under particular conditions. The determination of the sedimentation velocity was

- 5 classically performed in an analytical ultracentrifuge, but is typically now performed in a standard ultracentrifuge. See, Freifelder (1982) Physical Biochemistry (2d ed.), W.H. Freeman; and Cantor and Schimmel (1980) Biophysical Chemistry, parts 1-3, W.H. Freeman & Co., San Francisco; each of which is hereby incorporated herein by reference. As a crude determination, a sample containing a putatively soluble polypeptide is spun in a standard full sized ultracentrifuge at about 50K rpm for about 10 minutes, and soluble molecules will remain in the supernatant. A soluble particle or polypeptide will typically be less than about 30S, more typically less than about 15S, usually less than about 10S, more usually less than about 6S, and, in particular embodiments, preferably less than about 4S, and more preferably less than about 3S.

- A Langerin protein that specifically binds to or that is specifically immunoreactive with an antibody generated against a defined immunogen, such as an immunogen consisting of the Langerin protein of the invention is typically determined in an immunoassay. The immunoassay typically uses a polyclonal antiserum which was raised, e.g., to a Langerin protein of the invention. This antiserum is selected to have low crossreactivity against other potential Langerin family members, e.g., allelic variants of Langerin proteins preferably from the same species, and any such crossreactivity is removed by immunoabsorption prior to use in the immunoassay.

- To produce antisera for use in an immunoassay, the Langerin protein of the invention is isolated as described herein. For example, after determining the Langerin nucleic acid sequence using the methods described herein, a recombinant protein may be produced in a mammalian cell line. An appropriate host, e.g., an inbred strain of mice such as Balb/c, is immunized with

09787192.031501

the selected protein, typically using a standard adjuvant, such as Freund's adjuvant, and a standard mouse immunization protocol (see Harlow and Lane, *supra*). Alternatively, a synthetic peptide derived from a

5 Langerin nucleic acid sequence can be conjugated to a carrier protein and subsequently used as immunogen. Polyclonal sera are collected and titered against the immunogen protein in an immunoassay, e.g., a solid phase immunoassay with the immunogen immobilized on a solid

10 support. Polyclonal antisera with a titer of 10^4 or greater are selected and tested for their cross reactivity against other Langerin variants e.g., Langerin allelic variants, using a competitive binding immunoassay such as the one described in Harlow and Lane, *supra*, at

15 pages 570-573. Preferably at least two Langerin variants are used in this determination. These Langerin family members can be produced as recombinant proteins and isolated using standard molecular biology and protein chemistry techniques as described herein.

20 Immunoassays in the competitive binding format can be used for the crossreactivity determinations. For example, the Langerin protein of the invention can be immobilized to a solid support. Proteins added to the assay compete with the binding of the antisera to the

25 immobilized antigen. The ability of the above proteins to compete with the binding of the antisera to the immobilized protein is compared to a Langerin protein. The percent crossreactivity for the above proteins is calculated, using standard calculations. Those antisera

30 with less than 10% crossreactivity with each of the proteins listed above are selected and pooled. The cross-reacting antibodies are then removed from the pooled antisera by immunoabsorption with the above-listed proteins.

35 The immunoabsorbed and pooled antisera are then used in a competitive binding immunoassay as described above to compare a second Langerin protein to the immunogen protein (e.g., a Langerin allelic variant). To make this comparison, the two proteins are each assayed at a wide

09787192-031501

range of concentrations and the amount of each protein required to inhibit 50% of the binding of the antisera to the immobilized protein is determined. If the amount of the second protein required is less than twice the amount of the protein of the selected protein or proteins that is required, then the second protein is said to specifically bind to an antibody generated to the immunogen.

It is understood that the Langerin protein of the invention is not only the protein characterized herein, but also to other Langerin proteins that are allelic, non-allelic, or species variants. It is also understood that the terms include nonnatural mutations introduced by deliberate mutation using conventional recombinant technology such as single site mutation, or by excising short sections of DNA encoding the respective proteins, or by substituting new amino acids, or adding new amino acids to a Langerin sequence. Such minor alterations typically will substantially maintain the immunoidentity of the original molecule and/or its biological activity. Thus, these alterations include proteins that are specifically immunoreactive with a designated naturally occurring Langerin protein. The biological properties of altered proteins can be determined by expressing the protein in an appropriate cell line and measuring the appropriate effect, e.g., upon DC cells in vitro. Particular protein modifications considered minor include conservative substitution of amino acids with similar chemical properties. By aligning a protein optimally with the protein of the Langerins and by using conventional immunoassays as described herein to determine immunoidentity, one can determine the protein compositions of the invention.

IV. Physical Variants

This invention also encompasses proteins or peptides having substantial amino acid sequence homology with a natural mammalian Langerin protein or an antibody described above. The variants include species and

allelic variants. A person having ordinary skill in the art will recognize that much of the following discussion of variants will apply to variants of both the Langerin antigen and the antibodies which recognize it.

- 5 Amino acid sequence homology, or sequence identity, is determined by optimizing residue matches, if necessary, by introducing gaps as required. This changes when considering conservative substitutions as matches. Conservative substitutions typically include
- 10 substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine. Homologous amino acid sequences are typically intended to
- 15 include natural allelic and interspecies variations in each respective protein sequence. Typical homologous proteins or peptides will have from 25-100% homology (if gaps can be introduced), to 50-100% homology (if conservative substitutions are included) with the amino
- 20 acid sequence of the Langerin protein. Homology measures will be at least about 35%, generally at least 40%, more generally at least 45%, often at least 50%, more often at least 55%, typically at least 60%, more typically at least 65%, usually at least 70%, more usually at least
- 25 75%, preferably at least 80%, and more preferably at least 80%, and in particularly preferred embodiments, at least 85% or more. See also Needleham, et al. (1970) J. Mol. Biol. 48:443-453; Sankoff, et al. (1983) Chapter One in Time Warps, String Edits, and Macromolecules: The
- 30 Theory and Practice of Sequence Comparison Addison-Wesley, Reading, MA; and software packages from IntelliGenetics, Mountain View, CA; and the University of Wisconsin Genetics Computer Group, Madison, WI; each of which is incorporated herein by reference.
- 35 An isolated DNA, isolated as described below, encoding a Langerin protein can be readily modified by nucleotide substitutions, nucleotide deletions, nucleotide insertions, and inversions of nucleotide stretches. These modifications result in novel DNA

09787192.031501

sequences which encode these antigens, their derivatives, or proteins having similar physiological, immunogenic, or antigenic activity. These modified sequences can be used to produce mutant antigens or to enhance expression.

- 5 Enhanced expression may involve gene amplification, increased transcription, increased translation, and other mechanisms. Such mutant Langerin protein derivatives include predetermined or site-specific mutations of the respective protein or its fragments. "Mutant Langerin
10 protein" encompasses a polypeptide otherwise falling within the homology definition of the human Langerin protein as set forth above, but having an amino acid sequence which differs from that of Langerin protein as found in nature, whether by way of deletion,
15 substitution, or insertion. In particular, "site specific mutant Langerin protein" generally includes proteins having significant homology with a natural protein with properties described herein, and/or sharing various biological activities, e.g., antigenic or
20 immunogenic, with those sequences, and in preferred embodiments include, e.g., singly substituted natural forms of the proteins. Similar concepts apply to different Langerin proteins, particularly those found in various mammals, e.g., primates, including human. As
25 stated before, it is emphasized that the descriptions are generally meant to encompass all mammalian Langerin proteins.

- Although site specific mutation sites are predetermined, mutants need not be site specific.
- 30 Langerin protein mutagenesis can be conducted by making amino acid insertions or deletions. Substitutions, deletions, insertions, or any combinations may be generated to arrive at a final construct. Insertions include amino- or carboxy- terminal fusions. Random
35 mutagenesis can be conducted at a target codon and the expressed mutants can then be screened for the desired activity. Methods for making substitution mutations at predetermined sites in DNA having a known sequence are well known in the art, e.g., by M13 primer mutagenesis or

polymerase chain reaction (PCR) techniques. See also Sambrook, et al. (1989) and Ausubel, et al. (1987 and Supplements).

- 5 The mutations in the DNA normally should not place coding sequences out of reading frames and preferably will not create complementary regions that could hybridize to produce secondary mRNA structure such as loops or hairpins.

- 10 The present invention also provides recombinant proteins, e.g., heterologous fusion proteins using segments from these proteins. A heterologous fusion protein is a fusion of proteins or segments which are naturally not normally fused in the same manner. Thus, the fusion product of an immunoglobulin with a Langerin
15 polypeptide is a continuous protein molecule having sequences fused in a typical peptide linkage, typically made as a single translation product and exhibiting properties derived from each source peptide. A similar concept applies to heterologous nucleic acid sequences.

- 20 In addition, new constructs may be made from combining similar functional domains from other proteins. For example, antigen-binding or other segments may be "swapped" between different new fusion polypeptides or fragments. See, e.g., Cunningham, et al. (1989) Science
25 243:1330-1336; and O'Dowd, et al. (1988) J. Biol. Chem. 263:15985-15992, each of which is incorporated herein by reference. Thus, new chimeric polypeptides exhibiting new combinations of specificities will result from the functional linkage of biologically relevant domains and
30 other functional domains.

- The phosphoramidite method described by Beaucage and Carruthers (1981) Tetra. Letts. 22:1859-1862, will produce suitable synthetic DNA fragments. A double stranded fragment will often be obtained either by
35 synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence, e.g., PCR techniques.

V. Functional Variants

The blocking of physiological response mediated by Langerin proteins may result from the inhibition of binding of the antigen to its natural binding partner, e.g., through competitive inhibition. Thus, in vitro assays of the present invention will often use isolated protein, membranes from cells expressing a recombinant membrane associated Langerin protein, soluble fragments comprising binding segments, or fragments attached to solid phase substrates. These assays will also allow for the diagnostic determination of the effects of either binding segment mutations and modifications, or protein mutations and modifications, e.g., analogs. In particular, the Langerin is expressed on DC clones, but the antigen is lost after DC cell maturation.

This invention also contemplates the use of competitive drug screening assays, e.g., where neutralizing antibodies to antigen or binding partner fragments compete with a test compound for binding to the protein. In this manner, the antibodies can be used to detect the presence of any polypeptide which shares one or more antigenic binding sites of the protein and can also be used to occupy binding sites on the protein that might otherwise interact with a binding partner.

Additionally, neutralizing antibodies against the Langerin protein and soluble fragments of the antigen which contain a high affinity binding site, can be used to inhibit antigen function in cells or tissues, e.g., cells or tissues experiencing abnormal or undesired physiology. "Derivatives" of the Langerin antigens, and of antibodies, include amino acid sequence mutants, glycosylation variants, and covalent or aggregate conjugates with other chemical moieties. Covalent derivatives can be prepared by linkage of functionalities to groups which are found in the Langerin amino acid side chains or at the N- or C- termini, by means which are well known in the art. These derivatives can include, without limitation, aliphatic esters or amides of the carboxyl terminus, or of residues containing carboxyl

side chains, O-acyl derivatives of hydroxyl group-containing residues, and N-acyl derivatives of the amino terminal amino acid or amino-group containing residues, e.g., lysine or arginine. Acyl groups are selected from the group of alkyl-moieties including C3 to C18 normal alkyl, thereby forming alkanoyl aroyl species. Covalent attachment to carrier proteins may be important when immunogenic moieties are haptens.

In particular, glycosylation alterations are included, e.g., made by modifying the glycosylation patterns of a polypeptide during its synthesis and processing, or in further processing steps. Particularly preferred means for accomplishing this are by exposing the polypeptide to glycosylating enzymes derived from cells which normally provide such processing, e.g., mammalian glycosylation enzymes. Deglycosylation enzymes are also contemplated. Also embraced are versions of the same primary amino acid sequence which have other minor modifications, including phosphorylated amino acid residues, e.g., phosphotyrosine, phosphoserine, or phosphothreonine.

A major group of derivatives are covalent conjugates of the Langerin protein or fragments thereof with other proteins or polypeptides. These derivatives can be synthesized in recombinant culture such as N- or C-terminal fusions or by the use of agents known in the art for their usefulness in cross-linking proteins through reactive side groups. Preferred antigen derivatization sites with cross-linking agents are at free amino groups, carbohydrate moieties, and cysteine residues.

Fusion polypeptides between the Langerin proteins and other homologous or heterologous proteins are also provided. Homologous polypeptides may be fusions between different surface markers, resulting in, e.g., a hybrid protein exhibiting receptor binding specificity. Likewise, heterologous fusions may be constructed which would exhibit a combination of properties or activities of the derivative proteins. Typical examples are fusions of a reporter polypeptide, e.g., luciferase, with a

09787192-034501

segment or domain of amino acids of the antigen, e.g., a binding segment, so that the presence or location of the fused antigen may be easily determined. See, e.g., Dull, et al. U.S. Patent No. 4,859,609, which is hereby

- 5 incorporated herein by reference. Other gene fusion partners include bacterial β -galactosidase, trpE, Protein A, β -lactamase, alpha amylase, alcohol dehydrogenase, and yeast alpha mating factor. See, e.g., Godowski, et al. (1988) Science 241:812-816.

- 10 The phosphoramidite method described by Beaucage and Carruthers (1981) Tetra. Letts. 22:1859-1862, will produce suitable synthetic DNA fragments. A double stranded fragment will often be obtained either by synthesizing the complementary strand and annealing the
- 15 strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

Such polypeptides may also have amino acid residues which have been chemically modified by phosphorylation, sulfonation, biotinylation, or the addition or removal of

20 other moieties, particularly those which have molecular shapes similar to phosphate groups. In some embodiments, the modifications will be useful labeling reagents, or serve as purification targets, e.g., affinity ligands.

- 25 Fusion proteins will typically be made by either recombinant nucleic acid methods or by synthetic polypeptide methods. Techniques for nucleic acid manipulation and expression are described generally, for example, in Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual (Cur.. ed.), Vols. 1-3, Cold Spring Harbor Laboratory, which are incorporated herein by reference. Techniques for synthesis of polypeptides are described, e.g., in Merrifield (1963) J. Amer. Chem. Soc. 85:2149-2156; Merrifield (1986) Science 232: 341-347; and
- 30 Atherton, et al. (1989) Solid Phase Peptide Synthesis: A Practical Approach, IRL Press, Oxford; and Dawson, et al. (1994) Science 266:776-779; each of which is incorporated herein by reference.

This invention also contemplates the use of derivatives of the Langerin proteins other than variations in amino acid sequence or glycosylation. Such derivatives may involve covalent or aggregative association with chemical moieties. These derivatives generally fall into the three classes: (1) salts, (2) side chain and terminal residue covalent modifications, and (3) adsorption complexes, for example with cell membranes. Such covalent or aggregative derivatives are useful as immunogens, as reagents in immunoassays, or in purification methods such as for affinity purification of antigens or other binding proteins. For example, a Langerin antigen can be immobilized by covalent bonding to a solid support such as cyanogen bromide-activated Sepharose, by methods which are well known in the art, or adsorbed onto polyolefin surfaces, with or without glutaraldehyde cross-linking, for use in the assay or purification of anti-Langerin protein antibodies or other binding partner. Langerin antigens can also be labeled with a detectable group, for example radioiodinated by the chloramine T procedure, covalently bound to rare earth chelates, or conjugated to another fluorescent moiety for use in diagnostic assays. Purification of Langerin protein may be effected by immobilized antibodies or binding partners.

A solubilized Langerin antigen or fragment of this invention can be used as an immunogen for the production of antisera or antibodies specific for the protein or fragments thereof. The purified antigen can be used to screen monoclonal antibodies or binding fragments prepared by immunization with various forms of impure preparations containing the protein. In particular, antigen binding fragments of natural antibodies are often equivalent to the antibodies themselves. Purified Langerin protein can also be used as a reagent to detect any antibodies generated in response to the presence of elevated levels of the protein or cell fragments containing the antigen, both of which may be diagnostic of an abnormal or specific physiological or disease

09787192-031501

condition. Additionally, antigen fragments may also serve as immunogens to produce further antibodies of the present invention, as described immediately below. For example, this invention contemplates antibodies raised
5 against amino acid sequences from proteins having properties described in Table 1, or fragments of them. In particular, this invention contemplates antibodies having binding affinity to or being raised against specific fragments which are predicted to lie outside of
10 the lipid bilayer, e.g., either extracellular or intracellular domain structures.

The invention also provides means to isolate a group of related antigens displaying both distinctness and similarities in structure, expression, and function.
15 Elucidation of many of the physiological effects of the antigens will be greatly accelerated by the isolation and characterization of distinct species variants. In particular, the present invention provides useful probes for identifying additional homologous genetic entities in
20 different species.

Isolated genes will allow transformation of cells lacking expression of a corresponding Langerin protein, e.g., either species types or cells which lack corresponding antigens and should exhibit negative
25 background biological activity. Expression of transformed genes will allow isolation of antigenically pure cell lines, with defined or single specie variants. This approach will allow for more sensitive detection and discrimination of the physiological effects of Langerin
30 protein. Subcellular fragments, e.g., cytoplasts or membrane fragments, can also be isolated and used.

Dissection of the critical structural elements which effect the various physiological or differentiation functions provided by the proteins is possible using
35 standard techniques of modern molecular biology, particularly in comparing members of a related class. See, e.g., the homolog-scanning mutagenesis technique described in Cunningham, et al. (1989) Science 243:1339-1336; and approaches used in O'Dowd, et al. (1988) J.

Biol. Chem. 263:15985-15992; and Lechleiter, et al. (1990) EMBO J. 9:4381-4390; each of which is incorporated herein by reference.

In particular, functional domains or segments can be substituted between species variants or related proteins to determine what structural features are important in both binding partner affinity and specificity, as well as signal transduction. An array of different variants will be useful to screen for molecules exhibiting combined properties of interaction with different species variants of binding partners.

Antigen internalization may occur under certain circumstances, and interaction between intracellular components and "extracellular" segments of proteins involved in interactions may occur. The specific segments of interaction of Langerin with other intracellular components may be identified by mutagenesis or direct biochemical means, e.g., cross-linking or affinity methods. Structural analysis by crystallographic or other physical methods will also be applicable. Further investigation of the mechanism of biological function will include study of associated components which may be isolatable by affinity methods or by genetic means, e.g., complementation analysis of mutants.

Further study of the expression and control of Langerin will be pursued. The controlling elements associated with the antigens may exhibit differential developmental, tissue specific, or other expression patterns. Upstream or downstream genetic regions, e.g., control elements, are of interest.

Structural studies of the antigen will lead to design of new variants, particularly analogs exhibiting agonist or antagonist properties on binding partners. This can be combined with previously described screening methods to isolate variants exhibiting desired spectra of activities.

Expression in other cell types will often result in glycosylation differences in a particular antigen.

Various species variants may exhibit distinct functions based upon structural differences other than amino acid sequence. Differential modifications may be responsible for differential function, and elucidation of the effects are now made possible.

Thus, the present invention provides important reagents related to antigen-binding partner interaction. Although the foregoing description has focused primarily upon the human Langerin protein, those of skill in the art will immediately recognize that the invention encompasses other closely related antigens, e.g., other primate species or allelic variants, as well as variants and other members of the family.

VI. Nucleic Acids

The properties of natural Langerin protein disclosed herein allow for purification of the protein. This provides means to isolate nucleic acids encoding such proteins, and determination of the sequence provides a handle to other members of the Langerin family. Such nucleotide sequences and related reagents are useful in constructing a DNA clone useful for expressing Langerin protein, or, e.g., isolating a homologous gene from another natural source, including other members of the family. Typically, the sequences will be useful in isolating other genes, e.g., allelic variants or alternatively spliced isoforms, from human.

For example, a specific binding composition such as DCGM4 can be used for screening of an expression library made from a cell line which expresses a Langerin protein. The screening can be standard staining of surface expressed protein, or by panning. Screening of intracellular expression can also be performed by various staining or immunofluorescence procedures. The binding compositions could be used to affinity purify or sort out cells expressing the Langerin protein.

This invention contemplates use of isolated DNA or fragments to encode a biologically active Langerin protein or Langerin polypeptide. A nucleotide sequence

0978749-034501

encoding human langerin is shown in SEQ ID NO: 1. In addition, this invention covers isolated or recombinant DNA which encodes a biologically active Langerin protein or Langerin polypeptide and which is capable of

- 5 hybridizing under appropriate conditions with the DNA sequences. Said biologically active Langerin protein or Langerin polypeptide can be a full length antigen, or fragment thereof. Further, this invention covers the use of isolated or recombinant DNA, or fragments thereof,
10 which encode proteins which are homologous to a Langerin protein or which were isolated using cDNA encoding a Langerin protein as a probe. The isolated DNA can have the respective regulatory sequences in the 5' and 3' flanks, e.g., promoters, enhancers, poly-A addition
15 signals, and others.

- An "isolated" nucleic acid is a nucleic acid, e.g., an RNA, DNA, or a mixed polymer, which is substantially separated from other components which naturally accompany a native sequence, e.g., ribosomes, polymerases, and
20 flanking genomic sequences from the originating species. The term "isolated" nucleic acid embraces a nucleic acid sequence which has been removed from its naturally occurring environment, and it also includes recombinant or cloned DNA isolates and chemically synthesized analogs
25 or analogs biologically synthesized by heterologous systems. A substantially pure molecule includes isolated forms of the molecule. Alternatively, a purified species may be separated from host components from a recombinant expression system.

- 30 An isolated nucleic acid will generally be a homogeneous composition of molecules, but will, in some embodiments, contain minor heterogeneity. This heterogeneity is typically found at the polymer ends or portions not critical to a desired biological function or
35 activity.

A "recombinant" nucleic acid is defined either by its method of production or its structure. In reference to its method of production, e.g., a product made by a process, the process is use of recombinant nucleic acid

09787192.031501

techniques, e.g., involving human intervention in the nucleotide sequence, typically selection or production. Alternatively, it can be a nucleic acid made by generating a sequence comprising fusion of two fragments which are not naturally contiguous to each other, but is meant to exclude products of nature, e.g., naturally occurring mutants. Thus, for example, products made by transforming cells with any unnaturally occurring vector is encompassed, as are nucleic acids comprising sequence derived using any synthetic oligonucleotide process. Such is often done to replace a codon with a redundant codon encoding the same or a conservative amino acid, while typically introducing or removing a sequence recognition site. Alternatively, synthetic oligonucleotides are used to join together nucleic acid segments of desired functions to generate a single genetic entity comprising a desired combination of functions not found in the commonly available natural forms. Restriction enzyme recognition sites are often the target of such artificial manipulations, but other site specific targets, e.g., promoters, DNA replication sites, regulation sequences, control sequences, or other useful features may be incorporated by design. A similar concept is intended for a recombinant polypeptide, e.g., a fusion polypeptide. Specifically included are synthetic nucleic acids which, by genetic code redundancy, encode polypeptides similar to fragments of these antigens, and fusions of sequences from various different species variants.

A significant "fragment" in a nucleic acid context is a contiguous segment of at least about 17 nucleotides, generally at least 20 nucleotides, more generally at least 23 nucleotides, ordinarily at least 26 nucleotides, more ordinarily at least 29 nucleotides, often at least 32 nucleotides, more often at least 35 nucleotides, typically at least 38 nucleotides, more typically at least 41 nucleotides, usually at least 44 nucleotides, more usually at least 47 nucleotides, preferably at least 50 nucleotides, more preferably at least 53 nucleotides,

09787192-031501

and in particularly preferred embodiments will be at least 56 or more nucleotides.

- A DNA which codes for a Langerin protein will be particularly useful to identify genes, mRNA, and cDNA species which code for related or homologous proteins, as well as DNAs which code for homologous proteins. There should be homologues in other mammals, e.g., primates. Various Langerin proteins should be homologous and are encompassed herein. However, even proteins that have a more distant evolutionary relationship to the antigen can readily be isolated under appropriate conditions using these sequences if they are sufficiently homologous. Primate Langerin proteins are of particular interest.

- This invention further covers recombinant DNA molecules and fragments having a DNA sequence identical to or highly homologous to the isolated DNAs set forth herein. In particular, the sequences will often be operably linked to DNA segments which control transcription, translation, and DNA replication.
- Alternatively, recombinant clones derived from the genomic sequences, e.g., containing introns, will be useful for transgenic studies, including, e.g., transgenic cells and organisms, and for gene therapy. See, e.g., Goodnow (1992) "Transgenic Animals" in Roitt (ed.) Encyclopedia of Immunology Academic Press, San Diego, pp. 1502-1504; Travis (1992) Science 256:1392-1394; Kuhn, et al. (1991) Science 254:707-710; Capecchi (1989) Science 244:1288; Robertson (ed. 1987) Teratocarcinomas and Embryonic Stem Cells: A Practical Approach IRL Press, Oxford; Rosenberg (1992) J. Clinical Oncology 10:180-199; and Cournoyer and Caskey (1993) Ann. Rev. Immunol. 11:297-329; each of which is incorporated herein by reference.

- Homologous nucleic acid sequences, when compared, exhibit significant similarity. The standards for homology in nucleic acids are either measures for homology generally used in the art by sequence comparison or based upon hybridization conditions. The

09787492-034504

hybridization conditions are described in greater detail below.

- Substantial homology in the nucleic acid sequence comparison context means either that the segments, or
- 5 their complementary strands, when compared, are identical when optimally aligned, with appropriate nucleotide insertions or deletions, in at least about 50% of the nucleotides, generally at least 56%, more generally at least 59%, ordinarily at least 62%, more ordinarily at
- 10 least 65%, often at least 68%, more often at least 71%, typically at least 74%, more typically at least 77%, usually at least 80%, more usually at least about 85%, preferably at least about 90%, more preferably at least about 95 to 98% or more, and in particular embodiments,
- 15 as high at about 99% or more of the nucleotides. Alternatively, substantial homology exists when the segments will hybridize under selective hybridization conditions, to a strand, or its complement, typically using a sequence as described. Typically, selective
- 20 hybridization will occur when there is at least about 55% homology over a stretch of at least about 14 nucleotides, preferably at least about 65%, more preferably at least about 75%, and most preferably at least about 90%. See, Kanehisa (1984) Nuc. Acids Res. 12:203-213, which is
- 25 incorporated herein by reference. The length of homology comparison, as described, may be over longer stretches, and in certain embodiments will be over a stretch of at least about 17 nucleotides, usually at least about 20 nucleotides, more usually at least about 24 nucleotides,
- 30 typically at least about 28 nucleotides, more typically at least about 40 nucleotides, preferably at least about 50 nucleotides, and more preferably at least about 75 to 100 or more nucleotides.

- Stringent conditions, in referring to homology in
- 35 the hybridization context, will be stringent combined conditions of salt, temperature, organic solvents, and other parameters, typically those controlled in hybridization reactions. Stringent temperature conditions will usually include temperatures in excess of

00787192-134504

about 30° C, more usually in excess of about 37° C, typically in excess of about 45° C, more typically in excess of about 55° C, preferably in excess of about 65° C, and more preferably in excess of about 70° C.

- 5 Stringent salt conditions will ordinarily be less than about 1000 mM, usually less than about 500 mM, more usually less than about 400 mM, typically less than about 300 mM, preferably less than about 200 mM, and more preferably less than about 150 mM. However, the
- 10 combination of parameters is much more important than the measure of any single parameter. See, e.g., Wetmur and Davidson (1968) J. Mol. Biol. 31:349-370; Walker (ed. 1988) New Nucleic Acid Techniques Humana Press, Clifton, N.J.; Ross (ed. 1998) Nucleic Acid Hybridization; Wiley,
- 15 New York; or see, e.g., recent hybridization protocols on the Internet in "A Selection of Molecular Biology Protocols -- A list of sites with protocols," by Griffin (1996) Analytical Biochemistry 239:120-122.

20 VII. Making Langerin protein; Mimetics

- Langerin may be isolated from natural sources using standard methods of protein biochemistry. The DCGM4 antibody may be used to track the purification process, or in various immunoaffinity methods. Isolated protein
- 25 may be used as starting material for derivatization or modification. The protein may be used in native or denatured forms.

- DNA which encodes the Langerin protein or fragments thereof can be obtained by chemical synthesis, screening
- 30 cDNA libraries, or by screening genomic libraries prepared from a wide variety of cell lines or tissue samples. In particular, the DCGM4 antibody may be used to expression clone the nucleic acid encoding the protein antigen.

- 35 This DNA can be expressed in a wide variety of host cells for the synthesis of a full-length protein or fragments which can in turn, for example, be used to generate polyclonal or monoclonal antibodies; for binding studies; for construction and precision of modified

09757192-031501

molecules; and for structure/function studies. Each antigen or its fragments can be expressed in host cells that are transformed or transfected with appropriate expression vectors. These molecules can be substantially purified to be free of protein or cellular contaminants, other than those derived from the recombinant host, and therefore are particularly useful in pharmaceutical compositions when combined with a pharmaceutically acceptable carrier, buffer, and/or diluent. The antigen, or portions thereof, may be expressed as fusions with other proteins.

Expression vectors are typically self-replicating DNA or RNA constructs containing the desired antigen gene or its fragments, usually operably linked to suitable genetic control elements that are recognized in a suitable host cell. These control elements are capable of effecting expression within a suitable host. The specific type of control elements necessary to effect expression will depend upon the eventual host cell used. Generally, the genetic control elements can include a prokaryotic promoter system or a eukaryotic promoter expression control system, and typically include a transcriptional promoter, an optional operator to control the onset of transcription, transcription enhancers to elevate the level of mRNA expression, a sequence that encodes a suitable ribosome binding site, and sequences that terminate transcription and translation. Expression vectors also usually contain an origin of replication that allows the vector to replicate independently of the host cell.

The vectors of this invention contain DNA which encodes a Langerin protein, or a fragment thereof, preferably encoding a biologically active polypeptide. The sequence set forth in SEQ ID NO: 1 may advantageously be used to prepare Langerin protein. The DNA can be under the control of a viral promoter and can encode a selection marker. This invention further contemplates use of such expression vectors which are capable of expressing eukaryotic cDNA coding for a Langerin protein.

in a prokaryotic or eukaryotic host, where the vector is compatible with the host and where the eukaryotic cDNA coding for the antigen is inserted into the vector such that growth of the host containing the vector expresses the cDNA in question. Usually, expression vectors are designed for stable replication in their host cells or for amplification to greatly increase the total number of copies of the desirable gene per cell. It is not always necessary to require that an expression vector replicate in a host cell, e.g., it is possible to effect transient expression of the antigen or its fragments in various hosts using vectors that do not contain a replication origin that is recognized by the host cell. It is also possible to use vectors that cause integration of a Langerin gene or its fragments into the host DNA by recombination, or to integrate a promoter which controls expression of an endogenous gene e.g., see WO 96/29411 (incorporated herein by reference including all figures and drawings) describing such technology.

Vectors, as used herein, comprise plasmids, viruses, bacteriophage, integratable DNA fragments, and other vehicles which enable the integration of DNA fragments into the genome of the host. Expression vectors are specialized vectors which contain genetic control elements that effect expression of operably linked genes. Plasmids are the most commonly used form of vector but all other forms of vectors which serve an equivalent function and which are, or become, known in the art are suitable for use herein. See, e.g., Pouwels, et al. (1985 and Supplements) Cloning Vectors: A Laboratory Manual, Elsevier, N.Y.; and Rodriguez, et al. (eds. 1988) Vectors: A Survey of Molecular Cloning Vectors and Their Uses, Butterworth, Boston, MA; which are incorporated herein by reference.

Transformed cells include cells, preferably mammalian, that have been transformed or transfected with vectors containing a Langerin nucleic acid sequence, typically constructed using recombinant DNA techniques. Transformed host cells usually express the antigen or its

09787196-031501

09787193-034500

fragments, but for purposes of cloning, amplifying, and manipulating its DNA, do not need to express the protein. This invention further contemplates culturing transformed cells in a nutrient medium, thus permitting the protein to accumulate in the culture. The protein can be recovered, either from the culture or from the culture medium.

For purposes of this invention, DNA sequences are operably linked when they are functionally related to each other. For example, DNA for a presequence or secretory leader is operably linked to a polypeptide if it is expressed as a preprotein or participates in directing the polypeptide to the cell membrane or in secretion of the polypeptide. A promoter is operably linked to a coding sequence if it controls the transcription of the polypeptide; a ribosome binding site is operably linked to a coding sequence if it is positioned to permit translation. Usually, operably linked means contiguous and in reading frame, however, certain genetic elements such as repressor genes are not contiguously linked but still bind to operator sequences that in turn control expression.

Suitable host cells include prokaryotes, lower eukaryotes, and higher eukaryotes. Prokaryotes include both gram negative and gram positive organisms, e.g., *E. coli* and *B. subtilis*. Lower eukaryotes include yeasts, e.g., *S. cerevisiae* and *Pichia*, and species of the genus *Dictyostelium*. Higher eukaryotes include established tissue culture cell lines from animal cells, both of non-mammalian origin, e.g., insect cells, and birds, and of mammalian origin, e.g., human, primates, and rodents.

Prokaryotic host-vector systems include a wide variety of vectors for many different species. As used herein, *E. coli* and its vectors will be used generically to include equivalent vectors used in other prokaryotes. A representative vector for amplifying DNA is pBR322 or many of its derivatives. Vectors that can be used to express the Langerin proteins or its fragments include, but are not limited to, such vectors as those containing

the lac promoter (pUC-series); trp promoter (pBR322-trp); Ipp promoter (the pIN-series); lambda-pP or pR promoters (pOTS); or hybrid promoters such as ptac (pDR540). See Brosius, et al. (1988) "Expression Vectors Employing

- 5 Lambda-, trp-, lac-, and Ipp-derived Promoters", in Rodriguez and Denhardt (eds.) Vectors: A Survey of Molecular Cloning Vectors and Their Uses, Butterworth, Boston, Chapter 10, pp. 205-236, which is incorporated herein by reference.
- 10 Lower eukaryotes, e.g., yeasts and Dictyostelium, may be transformed with vectors encoding Langerin proteins. For purposes of this invention, the most common lower eukaryotic host is the baker's yeast, *Saccharomyces cerevisiae*. It will be used to generically
- 15 represent lower eukaryotes although a number of other strains and species are also available. Yeast vectors typically consist of a replication origin (unless of the integrating type), a selection gene, a promoter, DNA encoding the desired protein or its fragments, and
- 20 sequences for translation termination, polyadenylation, and transcription termination. Suitable expression vectors for yeast include such constitutive promoters as 3-phosphoglycerate kinase and various other glycolytic enzyme gene promoters or such inducible promoters as the alcohol dehydrogenase 2 promoter or metallothionine
- 25 promoter. Suitable vectors include derivatives of the following types: self-replicating low copy number (such as the YRp-series), self-replicating high copy number (such as the YE_p-series); integrating types (such as the
- 30 YIp-series), or mini-chromosomes (such as the YCp-series).

- Higher eukaryotic tissue culture cells are the preferred host cells for expression of the functionally active Langerin protein. In principle, any higher
- 35 eukaryotic tissue culture cell line is workable, e.g., insect baculovirus expression systems, whether from an invertebrate or vertebrate source. However, mammalian cells are preferred, in that the processing, both cotranslationally and posttranslationally.

09787192.031501

Transformation or transfection and propagation of such cells has become a routine procedure. Examples of useful cell lines include HeLa cells, Chinese hamster ovary (CHO) cell lines, baby rat kidney (BRK) cell lines, insect cell lines, bird cell lines, and monkey (COS) cell lines. Expression vectors for such cell lines usually include an origin of replication, a promoter, a translation initiation site, RNA splice sites (if genomic DNA is used), a polyadenylation site, and a transcription termination site. These vectors also usually contain a selection gene or amplification gene. Suitable expression vectors may be plasmids, viruses, or retroviruses carrying promoters derived, e.g., from such sources as adenovirus, SV40, parvoviruses, vaccinia virus, or cytomegalovirus. Representative examples of suitable expression vectors include pCDNA1; pCD, see Okayama, et al. (1985) Mol. Cell Biol. 5:1136-1142; pMCIneo Poly-A, see Thomas, et al. (1987) Cell 51:503-512; and a baculovirus vector such as pAC 373 or pAC 610.

It will often be desired to express a Langerin protein polypeptide in a system which provides a specific or defined glycosylation pattern. In this case, the usual pattern will be that provided naturally by the expression system. However, the pattern will be modifiable by exposing the polypeptide, e.g., an unglycosylated form, to appropriate glycosylating proteins introduced into a heterologous expression system. For example, the Langerin protein gene may be co-transformed with one or more genes encoding mammalian or other glycosylating enzymes. Using this approach, certain mammalian glycosylation patterns will be achievable or approximated in prokaryote or other cells.

The Langerin protein, or a fragment thereof, may be engineered to be phosphatidyl inositol (PI) linked to a cell membrane, but can be removed from membranes by treatment with a phosphatidyl inositol cleaving enzyme, e.g., phosphatidyl inositol phospholipase-C. This releases the antigen in a biologically active form, and allows purification by standard procedures of protein

09787192-031501

chemistry. See, e.g., Low (1989) Biochim. Biophys. Acta 988:427-454; Tse, et al. (1985) Science 230:1003-1008; and Brunner, et al. (1991) J. Cell Biol. 114:1275-1283.

- Fragments or derivatives of Langerin can be prepared
- 5 by conventional processes for synthesizing peptides. Solid phase and solution phase syntheses are both applicable to the foregoing processes. These include processes such as are described in Bodanszky (1993) Principles of Peptide Synthesis; Springer-Verlag;
 - 10 Bodanszky (1994) The Practice of Peptide Synthesis; Springer-Verlag; Jones (1992) Amino Acid and Peptide Synthesis; Oxford University Press; Atherton and Sheppard (1989) Solid Phase Peptide Synthesis: A Practical Approach; IRL Press New York; Stewart and Young (1984)
 - 15 Solid Phase Peptide Synthesis, Pierce Chemical Co., Rockford, IL; and chemical ligation, e.g., Dawson, et al. (1994) Science 266:776-779, a method of linking long synthetic peptides by a peptide bond; each of which is incorporated herein by reference. For example, an azide
 - 20 process, an acid chloride process, an acid anhydride process, a mixed anhydride process, an active ester process (for example, p-nitrophenyl ester, N-hydroxysuccinimide ester, or cyanomethyl ester), a carbodiimidazole process, an oxidative-reductive process,
 - 25 or a dicyclohexylcarbodiimide (DCCD)/additive process can be used. Solid phase and solution phase syntheses are both applicable to the foregoing processes.

- The Langerin protein, fragments, or derivatives are suitably prepared in accordance with the above processes
- 30 as typically employed in peptide synthesis, generally either by a so-called stepwise process which comprises condensing an amino acid to the terminal amino acid, one by one in sequence, or by coupling peptide fragments to the terminal amino acid. Amino groups that are not being
 - 35 used in the coupling reaction are typically protected to prevent coupling at an incorrect location.

If a solid phase synthesis is adopted, the C-terminal amino acid is bound to an insoluble carrier or support through its carboxyl group. The insoluble

00787192-001501

carrier is not particularly limited as long as it has a binding capability to a reactive carboxyl group.

Examples of such insoluble carriers include halomethyl resins, such as chloromethyl resin or bromomethyl resin, hydroxymethyl resins, phenol resins, tert-alkyloxycarbonyl-hydrazidated resins, and the like.

An amino group-protected amino acid is bound in sequence through condensation of its activated carboxyl group and the reactive amino group of the previously formed peptide or chain, to synthesize the peptide step by step. After synthesizing the complete sequence, the peptide is split off from the insoluble carrier to produce the peptide. This solid-phase approach is generally described by Merrifield, et al. (1963) in J. Am. Chem. Soc. 85:2149-2156, which is incorporated herein by reference.

The prepared protein and fragments thereof can be isolated and purified from the reaction mixture by means of peptide separation, for example, by extraction, precipitation, electrophoresis and various forms of chromatography, and the like. The Langerin of this invention can be obtained in varying degrees of purity depending upon its desired use. Purification can be accomplished by use of the protein purification techniques disclosed herein or by the use of the antibodies herein described in immunoabsorbant affinity chromatography. This immunoabsorbant affinity chromatography is carried out by first linking the antibodies to a solid support and then contacting the linked antibodies with solubilized lysates of appropriate source cells, lysates of other cells expressing the protein, or lysates or supernatants of cells producing Langerin protein as a result of DNA techniques, see below.

VIII. Utility

The present invention provides reagents which will find use in diagnostic applications as described elsewhere herein, e.g., in the general description for

00787133-001501

physiological or developmental abnormalities, or below in the description of kits. The antigen will be useful in both diagnostic methods, for isolating various cell types, e.g., Langerhans cells, or for forensic methods to
5 determine various species sources of biological samples.

This invention provides reagents with significant therapeutic value. Langerin (naturally occurring or recombinant), fragments thereof, muteins, and antibodies, along with compounds identified as having binding
10 affinity to Langerin or antibodies, find use in the treatment of conditions exhibiting abnormal expression of Langerin of its ligands or binding agents. Such abnormality will typically be manifested by immunological disorders. Additionally, this invention provides
15 therapeutic value in various diseases or disorders associated with abnormal expression or abnormal triggering of response to the ligand or binding agent. The Langerin ligands or binding agents are suspected to be involved in maturational development of DC. Langerin
20 may also be used as a receptor to target antigen to dendritic cells to elicit therapeutically relevant immune responses.

Recombinant Langerins, muteins, agonist or antagonist antibodies thereto, or antibodies can be
25 purified and then administered to a patient. These reagents can be combined for therapeutic use with additional active ingredients, e.g., in conventional pharmaceutically acceptable carriers or diluents, along with physiologically innocuous stabilizers and
30 excipients. These combinations can be sterile, e.g., filtered, and placed into dosage forms as by lyophilization in dosage vials or storage in stabilized aqueous preparations. This invention also contemplates the use of antibodies or binding fragments thereof which
35 are not complement binding.

Using Langerin or fragments thereof to screen for binding partner or for compounds having binding affinity to Langerin antigen can be performed including the isolation of associated compounds. Subsequent biological

09787192-034501

assays can then be utilized to determine if a putative ligand or binding agent can provide competitive binding, which can block intrinsic stimulating activity. Langerin fragments can be used as a blocker or antagonist in that it blocks the activity of ligand or binding agent. This invention further contemplates the therapeutic use of antibodies to Langerins as antagonists. This approach will be particularly useful with other Langerin protein species variants and other members of the family.

The quantities of reagents necessary for effective therapy will depend upon many different factors, including means of administration, target site, reagent physiological life, pharmacological life, physiological state of the patient, and other medicants administered. Thus, treatment dosages should be titrated to optimize safety and efficacy. Typically, dosages used in vitro may provide useful guidance in the amounts useful for in situ administration of these reagents. Animal testing of effective doses for treatment of particular disorders will provide further predictive indication of human dosage. The actual dosage of reagent, formulation or composition that modulates an immunological disorder depends on many factors, including the size and health of an organism, however, one of ordinary skill in the art can use teachings describing methods and techniques for determining clinical dosages. See, e.g., Spilker (1984) Guide to Clinical Studies and Developing Protocols, Raven Press Books, Ltd., New York, esp. pp. 7-13, 54-60; Spilker (1991) Guide to Clinical Trials, Raven Press, Ltd., New York, esp. pp. 93-101; Craig and Stitzel (eds. 1986) Modern Pharmacology, 2d ed., Little, Brown and Co., Boston, esp. pp. 127-33; Speight (ed. 1987) Avery's Drug Treatment: Principles and Practice of Clinical Pharmacology and Therapeutics, 3d ed., Williams and Wilkins, Baltimore, esp. pp. 50-56; Tallarida, et al. (1988) Principles of General Pharmacology, Springer-Verlag, New York, esp. pp. 18-20; Gilman, et al. (eds.) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, latest ed., Pergamon Press; Remington's

09787192-031504

09787192.031501

Pharmaceutical Sciences, latest ed., Mack Publishing Co., Easton, Penn.; and Rich, et al. (1998) Clinical Immunology: Principles and Practice Vols. I & II, Mosby, St. Louis, MO., each of which is hereby incorporated

5 herein by reference in its entirety including all drawings and figures. Methods for administration are discussed therein and below, e.g., for oral, intravenous, intraperitoneal, or intramuscular administration, transdermal diffusion, and others. Pharmaceutically

10 acceptable carriers will include water, saline, buffers, and other compounds described, e.g., in the Merck Index, Merck & Co., Rahway, New Jersey. Because of the likely high affinity binding, or turnover numbers, between a putative ligand or binding agent and its binding partner,

15 low dosages of these reagents would be initially expected to be effective. Thus, dosage ranges would ordinarily be expected to be in amounts lower than 1 mM concentrations, typically less than about 10 μ M concentrations, usually less than about 100 nM, preferably less than about 10 pM

20 (picomolar), and most preferably less than about 1 fM (femtomolar), with an appropriate carrier. Slow release formulations, or slow release apparatus will often be utilized for continuous administration.

Other abnormal developmental conditions are known in

25 the cell types shown to possess Langerin antigen, e.g., DC cells and epithelial cells of the tonsil. See Brew (ed.) The Merck Manual of Diagnosis and Therapy Merck & Co., Rahway, N.J.; and Thorn, et al. Harrison's Principles of Internal Medicine McGraw-Hill, N.Y. These

30 problems may be susceptible to prevention or treatment using compositions provided herein.

Langerins, fragments thereof, and antibodies or its fragments, antagonists, and agonists, may be administered directly to the host to be treated or, depending on the

35 size of the compounds, it may be desirable to conjugate them to carrier proteins such as ovalbumin or serum albumin prior to their administration. Therapeutic formulations may be administered in many conventional dosage formulations. While it is possible for the active

ingredient to be administered alone, it is preferable to present it as a pharmaceutical formulation. Formulations comprise at least one active ingredient, as defined above, together with one or more acceptable carriers thereof. Each carrier must be both pharmaceutically and physiologically acceptable in the sense of being compatible with the other ingredients and not injurious to the patient. Formulations include those suitable for oral, rectal, nasal, or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by methods well known in the art of pharmacy. See, e.g., Gilman, et al. (eds. 1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's Pharmaceutical Sciences, current ed., Mack Publishing Co., Easton, Penn.; Avis, et al. (eds. 1993) Pharmaceutical Dosage Forms: Parenteral Medications Dekker, NY; Lieberman, et al. (eds. 1990) Pharmaceutical Dosage Forms: Tablets Dekker, NY; and Lieberman, et al. (eds. 1990) Pharmaceutical Dosage Forms: Disperse Systems Dekker, NY. The therapy of this invention may be combined with or used in association with other therapeutic agents.

Both the naturally occurring and the recombinant forms of the Langerin specific antibodies and the Langerin proteins of this invention are particularly useful in kits and assay methods which are capable of screening compounds for binding activity. Several methods of automating assays have been developed in recent years so as to permit screening of tens of thousands of compounds in a short period. See, e.g., Fodor, et al. (1991) Science 251:767-773, which is incorporated herein by reference and which describes means for testing of binding affinity by a plurality of defined polymers synthesized on a solid substrate. The development of suitable assays can be greatly facilitated by the availability of large amounts of purified, soluble Langerin protein as provided by this invention.

09787192-033501

This invention is particularly useful for screening compounds by using recombinant antigen in any of a variety of drug screening techniques. The advantages of using a recombinant protein in screening for specific ligands or binding agents include: (a) improved renewable source of the antigen from a specific source; (b) potentially greater number of antigen molecules per cell giving better signal to noise ratio in assays; and (c) species variant specificity (theoretically giving greater biological and disease specificity). The purified protein may be tested in numerous assays, typically in vitro assays, which evaluate biologically relevant responses. See, e.g., Coligan Current Protocols in Immunology; Hood, et al. Immunology Benjamin/Cummings; Paul (ed. 1996) Fundamental Immunology 3d ed, Raven Press, NY; and Methods in Enzymology Academic Press. This will also be useful in screening for a ligand which binds a Langerin, e.g., from an interacting cell.

One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant DNA molecules expressing the Langerin antigens. Cells may be isolated which express an antigen in isolation from other functionally equivalent antigens. Such cells, either in viable or fixed form, can be used for standard protein-protein binding assays. See also, Parce, et al. (1989) Science 246:243-247; and Owicki, et al. (1990) Proc. Nat'l Acad. Sci. USA 87:4007-4011; which are incorporated herein by reference and describe sensitive methods to detect cellular responses.

Competitive assays are particularly useful, where the cells (source of Langerin protein) are contacted and incubated with a labeled binding partner or antibody having known binding affinity to the ligand, such as ^{125}I -antibody, and a test sample whose binding affinity to the binding composition is being measured. The bound and free labeled binding compositions are then separated to assess the degree of antigen binding. The amount of test compound bound is inversely proportional to the amount of labeled receptor binding to the known source.

- Numerous techniques can be used to separate bound from free antigen to assess the degree of binding. This separation step could typically involve a procedure such as adhesion to filters followed by washing, adhesion to plastic followed by washing, or centrifugation of the cell membranes. Viable cells could also be used to screen for the effects of drugs on Langerin protein mediated functions, e.g., second messenger levels, i.e., Ca^{++} ; cell proliferation; inositol phosphate pool changes; and others. Some detection methods allow for elimination of a separation step, e.g., a proximity sensitive detection system. Calcium sensitive dyes will be useful for detecting Ca^{++} levels, with a fluorimeter or a fluorescence cell sorting apparatus.
- Another method utilizes membranes from transformed eukaryotic or prokaryotic host cells as a source of Langerin protein. These cells are stably transformed with DNA vectors directing the expression of a membrane associated Langerin protein, e.g., an engineered membrane bound form. Essentially, the membranes would be prepared from the cells and used in a receptor/ligand type binding assay such as the competitive assay set forth above.
- Still another approach is to use solubilized, unpurified or solubilized, purified Langerin protein from transformed eukaryotic or prokaryotic host cells. This allows for a "molecular" binding assay with the advantages of increased specificity, the ability to automate, and high drug test throughput.
- Another technique for drug screening involves an approach which provides high throughput screening for compounds having suitable binding affinity to Langerin and is described in detail in Geysen, European Patent Application 84/03564, published on September 13, 1984, which is incorporated herein by reference. First, large numbers of different small peptide test compounds are synthesized on a solid substrate, e.g., plastic pins or some other appropriate surface, see Fodor, et al. (1991). Then all the pins are reacted with solubilized, unpurified or solubilized, purified Langerin binding

09787192-034504

composition, and washed. The next step involves detecting bound binding composition.

Rational drug design may also be based upon structural studies of the molecular shapes of the

5 Langerin protein and other effectors or analogs. Effectors may be other proteins which mediate other functions in response to antigen binding, or other proteins which normally interact with the antigen, e.g., Langerin ligand. One means for determining which sites

10 interact with specific other proteins is a physical structure determination, e.g., x-ray crystallography or 2 dimensional NMR techniques. These will provide guidance as to which amino acid residues form molecular contact

15 regions. For a detailed description of protein structural determination, see, e.g., Blundell and Johnson (1976) Protein Crystallography, Academic Press, New York, which is hereby incorporated herein by reference.

Purified Langerin protein can be coated directly

20 onto plates for use in the aforementioned drug screening techniques. However, non-neutralizing antibodies to these ligands can be used as capture antibodies to immobilize the respective ligand on the solid phase.

25 IX. Kits and Quantitation/Detection

Both naturally occurring and recombinant forms of the Langerin molecule of the invention are particularly useful in kits and assay methods. For example, these methods would also be applied to screening for binding

30 activity, e.g., ligands or binding agents for these proteins. Several methods of automating assays have been developed in recent years so as to permit screening of tens of thousands of compounds per year. See, e.g., a BIOMEK automated workstation, Beckman Instruments, Palo

35 Alto, California, and Fodor, et al. (1991) Science 251:767-773, which is incorporated herein by reference. The latter describes means for testing binding by an agent by a plurality of defined polymers synthesized on a solid substrate. The development of suitable assays to

09767192-034504

screen for a ligand or binding agent or agonist/antagonist homologous proteins can be greatly facilitated by the availability of large amounts of purified, soluble Langerin in an active state such as is provided by this invention.

Purified Langerin can be coated directly onto plates for use in the aforementioned ligand or binding agent screening techniques. However, non-neutralizing antibodies to these proteins can be used as capture antibodies to immobilize the respective Langerin protein on the solid phase, useful, e.g., in diagnostic uses.

This invention also contemplates use of Langerin, fragments thereof, peptides, and their fusion products in a variety of diagnostic kits and methods for detecting the presence of the Langerin protein or its ligand or binding agent. Alternatively, or additionally, antibodies against the Langerin molecules may be incorporated into the kits and methods. Typically the kit will have a compartment containing either a Langerin protein, fragment, peptide or gene segment or a reagent which recognizes one or the other of these. Typically, recognition reagents, in the case of a protein or fragment thereof, would be a receptor or antibody, or in the case of a gene segment, would usually be a hybridization probe.

A preferred kit for determining the concentration of Langerin in a sample would typically comprise a labeled compound, e.g., ligand, binding agent, or antibody, having known binding affinity for Langerin, a source of Langerin (naturally occurring or recombinant) as a positive control, and a means for separating the bound from free labeled compound, for example a solid phase for immobilizing the Langerin in the test sample. Compartments containing reagents, and instructions, will normally be provided.

This invention also contemplates use of Langerin proteins, fragments thereof, peptides, their fusion products, and binding compositions in a variety of diagnostic kits and methods for detecting the presence of

a binding composition. Typically the kit will have a compartment containing either a defined Langerin peptide or gene segment or a reagent which recognizes one or the other, e.g., antigen fragments or antibodies. See, e.g.,

5 Chen (ed.) (1987) Immunoassay: A Practical Guide Academic Press, Orlando, FL; Price and Newman (eds.) (1991) Principles and Practice of Immunoassay Stockton Press, New York; and Ngo (ed.) (1988) Nonisotopic Immunoassay Plenum Press, NY.

- 10 A kit for determining the binding affinity of a test compound to a Langerin protein would typically comprise a test compound; a labeled compound, for example an antibody having known binding affinity for the antigen; a source of Langerin protein (naturally occurring or
- 15 recombinant); and a means for separating bound from free labeled compound, such as a solid phase for immobilizing the antigen. Once compounds are screened, those having suitable binding affinity to the antigen can be evaluated in suitable biological assays, as are well known in the
- 20 art, to determine whether they exhibit similar biological activities to the natural antigen. The availability of recombinant Langerin protein polypeptides also provide well defined standards for calibrating such assays.

- One method for determining the concentration of
- 25 Langerin protein in a sample would typically comprise the steps of: (1) preparing membranes from a sample comprised of a membrane bound Langerin protein source; (2) washing the membranes and suspending them in a buffer; (3) solubilizing the antigen by incubating the membranes in a culture medium to which a suitable detergent has been
- 30 added; (4) adjusting the detergent concentration of the solubilized antigen; (5) contacting and incubating said dilution with radiolabeled antibody to form complexes; (6) recovering the complexes such as by filtration
- 35 through polyethyleneimine treated filters; and (7) measuring the radioactivity of the recovered complexes.

Antibodies, including antigen binding fragments, specific for the Langerin protein or fragments are useful in diagnostic applications to detect the presence of

09787192-031501

elevated levels of Langerin protein and/or its fragments. Such diagnostic assays can employ lysates, live cells, fixed cells, immunofluorescence, cell cultures, tissue samples, body fluids, and further can involve the

5 detection of antigens related to the protein in serum, or the like. Diagnostic assays may be homogeneous (without a separation step between free reagent and protein-protein complex) or heterogeneous (with a separation step). Various commercial assays exist, such as

10 radioimmunoassay (RIA), enzyme-linked immunosorbent assay (ELISA), enzyme immunoassay (EIA), enzyme-multiplied immunoassay technique (EMIT), substrate-labeled fluorescent immunoassay (SLFIA), and the like. For example, unlabeled antibodies can be employed by using a

15 second antibody which is labeled and which recognizes the antibody to a Langerin protein or to a particular fragment thereof. Similar assays have also been extensively discussed in the literature. See, e.g., Harlow and Lane (1988) Antibodies: A Laboratory Manual,

20 CSH, and Coligan (ed. 1991) and periodic supplements, Current Protocols In Immunology Greene/Wiley, New York.

Anti-idiotypic antibodies may have similar use to diagnose presence of antibodies against a Langerin protein, as such may be diagnostic of various abnormal

25 states. For example, overproduction of Langerin protein may result in production of various immunological reactions which may be diagnostic of abnormal physiological states associated with Langerhans cells (e.g., atopic dermatitis, eosinophilic granuloma,

30 histiocytosis, systemic sclerosis and Langerhans cell granulomatosis). Also, see, e.g., Rich, et al. (1998) Clinical Immunology: Principles and Practice Vols. I & II, Mosby, St. Louis, Mo.

Frequently, the reagents for diagnostic assays are

35 supplied in kits, so as to optimize the sensitivity of the assay. For the subject invention, depending upon the nature of the assay, the protocol, and the label, either labeled or unlabeled antibody, or labeled Langerin protein is provided. This is usually in conjunction with

09787193-031501

09787162-031404

other additives, such as buffers, stabilizers, materials necessary for signal production such as substrates for enzymes, and the like. Preferably, the kit will also contain instructions for proper use and disposal of the contents after use. Typically the kit has compartments for each useful reagent. Desirably, the reagents are provided as a dry lyophilized powder, where the reagents may be reconstituted in an aqueous medium providing appropriate concentrations of reagents for performing the assay.

Many of the aforementioned constituents of the drug screening and the diagnostic assays may be used without modification or may be modified in a variety of ways. For example, labeling may be achieved by covalently or non-covalently joining a moiety which directly or indirectly provides a detectable signal. In these assays, the antigen, test compound, Langerin protein, or antibodies thereto can be labeled either directly or indirectly. Possibilities for direct labeling include label groups: radiolabels such as ^{125}I , enzymes (U.S. Pat. No. 3,645,090) such as peroxidase and alkaline phosphatase, and fluorescent labels (U.S. Pat. No. 3,940,475) capable of monitoring the change in fluorescence intensity, wavelength shift, or fluorescence polarization. Both of the patents are incorporated herein by reference. Possibilities for indirect labeling include biotinylation of one constituent followed by binding to avidin coupled to one of the above label groups.

There are also numerous methods of separating the bound from the free antigen, or alternatively the bound from the free test compound. The Langerin protein can be immobilized on various matrices followed by washing. Suitable matrices include plastic such as an ELISA plate, filters, and beads. Methods of immobilizing the Langerin protein to a matrix include, without limitation, direct adhesion to plastic, use of a capture antibody, chemical coupling, and biotin-avidin. The last step in this approach involves the precipitation of protein-protein

complex by any of several methods including those utilizing, e.g., an organic solvent such as polyethylene glycol or a salt such as ammonium sulfate. Other suitable separation techniques include, without
5 limitation, the fluorescein antibody magnetizable particle method described in Rattle, et al. (1984) Clin. Chem. 30:1457-1461, and the double antibody magnetic particle separation as described in U.S. Pat. No. 4,659,678, each of which is incorporated herein by
10 reference.

The methods for linking proteins or their fragments to the various labels have been extensively reported in the literature and do not require detailed discussion here. Many of the techniques involve the use of
15 activated carboxyl groups either through the use of carbodiimide or active esters to form peptide bonds, the formation of thioethers by reaction of a mercapto group with an activated halogen such as chloroacetyl, or an activated olefin such as maleimide, for linkage, or the
20 like. Fusion proteins will also find use in these applications.

Another diagnostic aspect of this invention involves use of oligonucleotide or polynucleotide sequences taken from the sequence of a Langerin protein. These sequences can
25 be used as probes for detecting levels of antigen message in samples from patients suspected of having an abnormal condition, e.g., an immunological disorder. The preparation of both RNA and DNA nucleotide sequences, the labeling of the sequences, and the preferred size of the
30 sequences has received ample description and discussion in the literature. Normally an oligonucleotide probe should have at least about 14 nucleotides, usually at least about 18 nucleotides, and the polynucleotide probes may be up to several kilobases. Various labels may be
35 employed, most commonly radionuclides, particularly ^{32}P . However, other techniques may also be employed, such as using biotin modified nucleotides for introduction into a polynucleotide. The biotin then serves as the site for binding to avidin or antibodies, which may be labeled

with a wide variety of labels, such as radionuclides, fluorescers, enzymes, or the like.

- Alternatively, antibodies may be employed which can recognize specific duplexes, including DNA duplexes, RNA duplexes, DNA-RNA hybrid duplexes, or DNA-protein duplexes. The antibodies in turn may be labeled and the assay carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.
- The use of probes to the novel anti-sense RNA may be carried out in any conventional techniques such as nucleic acid hybridization, plus and minus screening, recombinational probing, hybrid released translation (HRT), and hybrid arrested translation (HART). This also includes amplification techniques such as polymerase chain reaction (PCR).

- Diagnostic kits which also test for the qualitative or quantitative presence of other markers are also contemplated. Diagnosis or prognosis may depend on the combination of multiple indications used as markers. Thus, kits may test for combinations of markers. See, e.g., Viallet, et al. (1989) Progress in Growth Factor Res. 1:89-97.

25 X. Isolating Langerin Specific Binding Partners

- The description of the Langerin protein herein provides means to identify Langerin ligands or binding agents, as described above. Such a ligand or binding agent should bind specifically, e.g., selectively, to the Langerin or fragment thereof with reasonably high affinity. Typical ligand or binding agent binding constants will be at least about 30 mM, e.g., generally at least about 3 mM, more generally at least about 300 μ M, typically at least about 30 μ M, 3 μ M, 300 nM, 30 nM, etc. Various constructs are made available which allow either labeling of Langerin to detect its ligand or binding agent. For example, directly labeling Langerin, fusing onto it markers for secondary labeling, e.g., FLAG or other epitope tags, etc., will allow detection of a

005787492-001504

binding agent or ligand. This can be histological, as an affinity method for biochemical purification, or labeling or selection in an expression cloning approach. A two-hybrid selection system may also be applied making appropriate constructs with the available Langerin sequences. See, e.g., Fields and Song (1989) Nature 340:245-246.

The Langerin protein should interact with a ligand based, e.g., upon its similarity in structure and function to other cell markers exhibiting developmental and cell type specificity of expression. Methods to isolate a ligand are made available by the ability to make purified Langerin for screening programs. Soluble or other constructs using the Langerin sequences provided herein will allow for screening or isolation of Langerin specific ligands.

Generally, descriptions of Langerins will be analogously applicable to individual specific embodiments directed to Langerin and/or Langerin reagents and compositions.

Without further elaboration, it is believed that a person of ordinary skill in the art can, using the preceding description, utilize the present invention to its fullest extent. The following examples are put forth solely for the purpose of illustration as to make and use the present invention, and are not intended, nor should they be construed, to limit the scope of what the inventors regard as their invention. Unless indicated otherwise below, parts are parts by weight, molecular weight is weight average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

EXAMPLES

General Methods

Some of the standard methods are described or referenced, e.g., in Maniatis, et al. (1982) Molecular Cloning, A Laboratory Manual Cold Spring Harbor Laboratory, Cold Spring Harbor Press; Sambrook, et al.

- (1989) Molecular Cloning: A Laboratory Manual (2d ed.), vols 1-3, CSH Press, NY; Ausubel, et al., Biology, Greene Publishing Associates, Brooklyn, NY; or Ausubel, et al. (1987 and Supplements) Current Protocols in Molecular
- 5 Biology Greene/Wiley, New York; Innis, et al. (eds. 1990) PCR Protocols: A Guide to Methods and Applications Academic Press, N.Y.; all of which are each incorporated herein by reference. Methods for protein purification include such methods as ammonium sulfate precipitation,
- 10 column chromatography, electrophoresis, centrifugation, crystallization, and others. See, e.g., Ausubel, et al. (1987 and periodic supplements); Deutscher (1990) "Guide to Protein Purification," Methods in Enzymology vol. 182, and other volumes in this series; Coligan, et al. (1995
- 15 and supplements) Current Protocols in Protein Science John Wiley and Sons, New York, NY; Matsudaira (ed. 1993) A Practical Guide to Protein and Peptide Purification for Microsequencing, Academic Press, San Diego, CA; and manufacturer's literature on use of protein purification
- 20 products, e.g., Pharmacia, Piscataway, NJ, or Bio-Rad, Richmond, CA. Combination with recombinant techniques allow fusion to appropriate segments, e.g., to a FLAG sequence or an equivalent which can be fused via a protease-removable sequence. See, e.g., Hochuli (1989)
- 25 Chemische Industrie 12:69-70; Hochuli (1990) "Purification of Recombinant Proteins with Metal Chelate Absorbent" in Setlow (ed.) Genetic Engineering, Principle and Methods 12:87-98, Plenum Press, N.Y.; and Crowe, et al. (1992) QIAexpress: The High Level Expression &
- 30 Protein Purification System QUIAGEN, Inc., Chatsworth, CA: which are incorporated herein by reference.
- FACS analyses are described in Melamed, et al. (1990) Flow Cytometry and Sorting Wiley-Liss, Inc., New York, NY; Shapiro (1988) Practical Flow Cytometry Liss, New York, NY; and Robinson, et al. (1993) Handbook of Flow Cytometry Methods Wiley-Liss, New York, NY.

09787192-031501

Example 1
Dendritic Cell Clones

Collection and Purification of cord blood CD34⁺ HPC

- 5 Umbilical cord blood samples were obtained according to standard institutional guidelines for human samples. Cells bearing CD34 antigen were isolated from mononuclear fractions by positive selection with Minimacs separation columns (Miltenyi Biotec, Bergish Gladbach, Germany),
10 using an anti-CD34 mAb (Immu-133.3, Immunotech, Marseille, France) and goat anti-mouse IgG-coated microbeads (Miltenyi Biotec). In all experiments, isolated cells were 80-99% CD34⁺ as judged by staining with an anti-CD34 mAb.

- 15 Hematopoietic factors

- Recombinant human (rh) GM-CSF (specific activity: 2×10^6 U/mg; Schering-Plough Research Institute, Kenilworth, NJ) was used at 100 ng/ml (200 U/ml); rhTNF- α
20 (specific activity: 2×10^7 U/mg; Genzyme, Boston, MA) was used at 2.5 ng/ml (50 U/ml); rhSCF (specific activity: 4×10^5 U/mg; R&D, Abington, UK) was used at 25 ng/ml; rhIL-4 (specific activity: 10^7 U/mg; Schering-Plough Research Institute) was used at 5 ng/ml; and
25 rhTGF- β 1 (R&D) was used at 1 ng/ml.

Dendritic cell generation from CD34⁺ HPC

- Cultures of CD34⁺ HPC were established as described in Caux, et al. (1992) Nature 392:258-261, in the
30 presence of GM-CSF, SCF, and TNF- α , in endotoxin-free medium consisting of RPMI 1640 (Gibco BRL, Gaithersburg, MD) supplemented with 10% heat-inactivated fetal bovine serum (FBS; Flow Laboratories, Irvine, UK), 10 mM Hepes, 2 mM L-glutamine, 5×10^{-5} M 2-mercaptoethanol and
35 gentamicin (80 μ g/ml) (referred to as complete medium).

- CD34⁺ cells were seeded in 25 to 75 cm² culture flasks (Corning, New York; NY) at 2×10^4 cells/ml. Optimal conditions were maintained by splitting cultures at day 4 with medium containing fresh GM-CSF and TNF- α .
40 In some experiments, TNF- α was replaced at day 7 by

- TGF- β . In other experiments, DC were activated at day 9 or day 12 with murine Ltk⁻ fibroblastic L cells stably transfected with the human CD40 ligand (L) gene (cell line established by Dr. C. van Kooten (van Kooten, et al. (1994) Eur J Immunol 24:787-92). Briefly, 10⁵ irradiated CD40L L cells (7,500 rads) were seeded together with 5 10⁵ CD34-derived DC in presence of GM-CSF.

Isolation of epidermal cells

- Epidermal cell suspensions were obtained from normal skin of patients undergoing reconstructive plastic surgery of the breast. Skin was split-cut with a keratome set and the dermo-epidermal slices treated for 18 h at 4° C with 0.05% trypsin (Sigma) in Hank's balanced salt solution without Ca²⁺ and Mg²⁺ (Seromed, Biochrom KG, Berlin, FRG). The epidermis was detached from the dermis with fine forceps. Epidermal sheet cell suspensions were obtained by subsequent tissue dislocation and filtration through sterile gaze.
- Enrichment of Langerhans cells was obtained by density gradient centrifugation on Lymphoprep (Nycomed Pharma, Oslo, Norway).

Example 2

- Antibodies and Flow Cytometry

- For single cell staining, cells were labeled using the following mAbs: anti-E-cadherin (SHE 79.7; Takara, Shiga, Japan), anti-MHC class II (HLA-DR) (Becton Dickinson), and anti-Lag, all revealed by FITC-conjugated goat anti-mouse immunoglobulin (Dako, Glostrup, Denmark). For double staining, cells were labeled with mAb DCGM4 revealed by phycoerythrin(PE)-conjugated goat anti-mouse immunoglobulin (Dako), and after saturation in 5% mouse serum, with FITC-conjugated anti-CD1a (Ortho, Raritan, NJ). Negative controls were performed with unrelated murine mAbs. Fluorescence was determined with a FACSCAN flow-cytometer (Becton Dickinson). For intracytoplasmic phenotyping, cells were stained in PBS, 0.3% Saponin (Sigma) and 5% BSA, using the same procedure.

09787192-031504

Example 3
Biochemistry

Proteins were extracted from CD34-derived DC
5 supplemented with TGF- β by addition, to a frozen pellet
of 100 μ l/10⁷ cells, of 50 mM Tris-HCL pH 8 buffer with
150 mM NaCl, 5 mM EDTA, 1% Triton X100 and protease
inhibitor (complete Mini, Boeringer Mannheim). After 1 h
at 4° C, samples were centrifuged to remove cellular
10 debris. Supernatants were then incubated 1 h at 4° C
with DCGM4 covalently linked to Dynabeads M-450 Sheep
anti-mouse magnetic beads (Dyna, Oslo, Norway). Beads
were washed with extraction buffer by using a Dynal
magnetic particle-concentrator and boiled in the presence
15 of 50 μ l SDS-PAGE sample buffer or resuspended in 100 μ l
of 0.5 M Glycine, 0.15 M NaCl pH 2.3 for 4 minutes.
Then, supernatant was neutralized with 3.5 μ l of
saturated Tris solution. SDS-PAGE analysis was performed
with a PhastSystem in a 10-15% gradient gel (Pharmacia
20 Biotech), and gels were stained with Coomassie R250. 2-D
analysis was performed on a Multiphor II flat bed system
with Immobiline DryStrip pH 3-10 and Excelgel SDS 8-18%
for the second dimension (Pharmacia Biotech), and gels
were silver stained. Dialyzed samples of proteins eluted
25 from IgG linked to Dynabeads were digested with N-
glycosidase F (Boehringer-Mannheim) at 37° C overnight.
Five microliters of original samples or 10 microliters of
digested samples were deposited on nitrocellulose, and
treated with DIG glycan detection kit (Boehringer-
30 Mannheim).

Example 4
Purification of Langerin

35 DCGM4 expressing cells are used for large scale
protein extraction. The protein is gently solubilized,
and the resulting Langerin is purified using standard
methods of protein purification. Chromatographic methods
are used, and immunoaffinity techniques can be applied.

The Langerin protein is followed by SDS PAGE and/or immunoassays. Diagnostic methods are used to ensure that the protein is substantially pure.

- Purified protein is used for protein
- 5 microsequencing. See, e.g., Matsudaira (ed. 1993) A Practical Guide to Protein and Peptide Purification for Microsequencing, Academic Press, San Diego, CA. Sequence data is used to search sequence databases, e.g., GENBANK, to find natural genes encoding the Langerin.
 - 10 Alternatively, the sequence data is useful for isolating a nucleic acid encoding Langerin using, e.g., degenerate PCR primers, etc.

- Protein will also be used to raise additional antibodies. Such antibodies may be polyclonal or
- 15 monoclonal. The protein can be used to assay and determine titers and affinity. Standard methods of immunization are available, as described above.

Example 5

- 20 Internalization Assay

- CD34-derived DC supplemented with TGF- β were generated as detailed above and internalization was performed as described (Cella, et al. (1997) J. Exp. Med.
- 25 185:1743-51). One aliquot of cells was fixed with RPMI, 0.1% glutaraldehyde for 5 min. at room temperature and another aliquot was used without fixation. Both samples were stained with mAb DCGM4 or mAb DCGM1 for 40 min. on ice, and incubated with biotin-labeled F(ab')₂ goat anti-
 - 30 mouse IgG (Jackson ImmunoResearch Laboratories, PA) for 1 h on ice. Cells were then placed in a 37°C water-bath for various time periods, cooled on ice and stained with PE-conjugated streptavidin (Becton Dickinson). After washing, cells were analyzed by FACS. The measure of
 - 35 internalization is given by the percentage decrease of cell-surface median fluorescence intensity (MFI) as compared to control samples kept at 4° C. The percentage decrease of MFI observed in fixed cells was taken as measure of the off-rate of the antibody at 37° C. Linear

regression analysis of the plots of \log_{10} (percentage of median fluorescence) vs. time was performed and rates of ingestion (k) and half-life ($t_{1/2}$) of membrane-bound complexes were calculated from the slope (m) of the resulting straight line using the relationships $k(\%/min) = -2,303m \times 100$, and $t_{1/2} (min) = \log 2/m$ (Leslie, EJI 1980). In this mAb DCGM1 which recognizes the macrophage mannose receptor was generated by Applicant, and used as positive control for receptor-mediated endocytosis.

Example 6
Generation and Characterization of
DCGM4 Monoclonal Antibodies

BALB/c mice (Iffa Credo, Les Oncins, France) were immunized with three intraperitoneal injections of CD34-derived DC (10^6 cells) with Freund's adjuvant (Sigma Chemical Co., St. Louis, MO). Three days after the final injection, splenocytes were fused with the murine myeloma cell line SP2, using polyethylene glycol-1000 (Sigma). Hybrid cells were placed in 96-well Falcon tissue culture plates (Falcon, Lincoln Park, NJ) and fed with DMEM F12 (Life Technologies, Gaithersburg, MD) supplemented with streptomycin (100 μ g/ml), penicillin (100 U/ml), glutamine (2 mM), 10% horse serum (Life Technologies), 1% culture medium additive (CRTS, Lyon, France), 10^{-5} M azaserine (Sigma), and 5×10^{-5} M hypoxanthine. Supernatants were screened for reactivity with CD34-derived DC and three unrelated cell types, namely peripheral blood polynuclear cells, T lymphocytes activated with PHA, and the myeloid cell line KG1 (ATCC; Rockville, MD). Selected hybridomas were cloned by limiting dilution and ascites were produced in BALB/c mice. MAb DCGM4 was purified by anion-exchange chromatography on DEAE A50 (Pharmacia Biotech, Uppsala, Sweden) and coupled with fluorescein and biotin using standard procedures. Ig isotype was determined by ELISA using a mouse hybridoma subtyping kit (Boehringer-Mannheim, Mannheim, Germany).

09787192-031501

Example 7
Immunohistology

Microscope slides of acetone-fixed cryocut tissue sections or cell cytospin preparations were incubated with mAbs for 60 min., and subsequently with biotinylated sheep anti-mouse Ig (The Binding Site, Birmingham, UK) for 30 min. Following incubation with streptavidin coupled to alkaline phosphatase (Biosource, CA, USA) for 30 min., enzyme activity was developed using Fast Red substrate (Dako). Double staining, with mouse IgG1 antibody DCGM4 and IgG2b anti-CD1a (Immunotech) were revealed by sheep anti-mouse IgG1 (The Binding Site) followed by mouse anti-alkaline phosphatase-alkaline phosphatase complexes (Dako) (APAAP technique), and biotinylated sheep anti-mouse IgG2b (The Binding Site) followed by ExtrAvidin-peroxydase (Sigma). The binding of goat anti-sIgD-biotin and DCGM4-biotin (for double staining with Lag) were directly revealed by ExtrAvidin-peroxidase. Alkaline phosphatase activity and peroxidase activity were respectively demonstrated using Fast Blue substrate (Sigma) and 3-amino-ethylcarbazole (Sigma).

Example 8
Electron Microscopy

Langerhans cell-enriched epidermal cell suspensions were incubated with control mouse IgG1 (Sigma), anti-CD1a (DMC1 mAb; REF), or DCGM4 for 1 h at 4°C. After washing, cells were incubated with a goat anti-mouse IgG conjugated with colloidal gold particles of 5 nM (GAM-nM) (Amersham, address, France) for 30 min. at 4°C. Cells were either fixed immediately for 18 hours with 2% glutaraldehyde in cacodylate buffer, followed by washing for at least 24 h in cacodylate buffer with sucrose, or warmed up to 37°C or room temperature before fixation. Samples were post-fixed for 1 hour with 1% osmium in cacodylate buffer with sucrose, dehydrated, and embedded

09787193-001304

- in Epoxy resin. Ultrathin sections were post-stained with uranyl acetate and lead citrate, and examined on a JEOL 1200 EX electron microscope (CMEABG, Université de Lyon, Lyon, France). A quantitative evaluation of cell-surface antigen density was performed according to Lafferty, et al. (1981) J. Histochem Cytochem 29:49-56. The number of gold granules bound along the cell membrane was counted and total circumference of the cell was measured on micrographs with a minimop morphometric analyzer (Zeiss).
- Results were expressed as number of gold granules per 100 μm of cell membrane. For each group of cells, counts were obtained from at least 20-30 cell sections. Mean and standard deviation were subsequently determined.

Example 9

Confocal Microscopy

- Intracellular immunofluorescence staining was performed as previously described by Winzler, et al. (1997) J. Exp. Med. 185:317-28. Cells on polylysine coated coverslips and fixed for 15 min. with 4% paraformaldehyde, were washed in 10 mM glycine, then permeabilized with 0.5% saponin, 0.2% BSA for 30 min. Coverslips were incubated for 30 min. at room temperature with anti-LAMP-1 (PharMingen, San Diego, CA), anti-HLA-DR (Becton Dickinson) or anti-Lag (Kashishara, et al. J Invest Dermatol 87:602-607) at a final concentration of 5 $\mu\text{g}/\text{ml}$ in permeabilization medium. After three washes, cells were incubated for 30 min. with secondary labeled antibody (donkey anti-mouse coupled to Texas-red (Vector Laboratories, Burlingame, CA), washed, and incubated with mouse preimmune serum for 30 min., washed again, post-fixed with 2% paraformaldehyde, and finally incubated for 30 min. with mAb DCGM4 coupled to fluorescein (FITC).
- After washing, coverslips were mounted onto glass slides with fluoromount (Southern Biotechnology Associates Inc., Birmingham, AL). Confocal microscopy was performed using Confocal Laser Scanning Microscope TCS 4D (Leica Lasertechnik GmbH, Heidelberg, Germany) interfaced with

an argon/krypton ion laser and with fluorescence filters and detectors allowing to simultaneously record FITC and Texas-red markers Rovere, et al. (1998) Proc. Nat. Acad. Sci. USA 95:1067-1072.

5

Example 10

Selection of Monoclonal Antibody DCGM4 Reactive Against Langerin

- 10 Supernatants from 854 hybridomas were screened for reactivity on DC obtained from CD34⁺ HPC cultured 12 days in GM-CSF and TNF- α . In parallel, the supernatants were assayed for reactivity on three unrelated cell types, namely peripheral blood polynuclear cells, T lymphocytes
- 15 activated with PHA and the myeloid cell line KG1. Supernatant from one hybridoma, designated DCGM4 was found to react only with a minor subset of DC and not with the other cell types of the differential screening.

- The hybridoma was cloned by limiting dilution. The
- 20 antibody was produced in ascites and subsequently purified using DEAE chromatography. MAb DCGM4 was found to be of IgG1/ κ isotype as determined by ELISA. As the observed reactivity restricted to a DC subset appeared of particular interest, mAb DCGM4 was selected for further
- 25 studies. Finally, early experiments indicating that LC stained positive, we termed Langerin the antigen recognized by mAb DCGM4.

Example 11

- 30 Langerin is Selectively Expressed
on Langerhans-Type Immature Dendritic Cells

- CD34⁺ HPC cultured with a combination of GM-CSF and TNF- α for 12 days differentiate into CD1a⁺ DC (Caux, et
- 35 al. (1996) J Exp Med 184:695-706). The expression of Langerin during such cultures was investigated. Langerin is expressed by a subset of CD1a⁺ dendritic cells derived from cord blood CD34⁺ HPC. Kinetics of CD1a and Langerin expression during culture of CD34⁺ HPC in GM-CSF plus

09787192-031501
105760-25787660

TNF- α was determined at various time points, cells were recovered and double-labeled using anti-CD1a-FITC and DCGM4 plus anti-mouse IgG-PE. The results are representative of 5 experiments.

5 No staining was detected at day 0 or day 6, indicating that CD34⁺ HPC and their immediate progeny do not express Langerin. The antigen appeared at day 7, on a small subset of the CD1a⁺ cells. Between day 7 to day 12, Langerin expression reached a maximum, staining
10 between 15 to 35% of CD1a⁺ cells. The dendritic nature of the Langerin-expressing cells in the cultures was confirmed on cytopsin preparations of DCGM4⁺ FACS-sorted cells.

 Caux, et al. (1996) J Exp Med 184:695-706 have
15 further shown that CD34⁺ HPC differentiate along two independent pathways from distinct precursor subsets, identified by mutually exclusive expression of CD1a and CD14 at early time points during the culture (day 5-7). When such precursors were separated by FACS-sorting at
20 day 6 and cultured with GM-CSF and TNF- α for 6 more days, Langerin was found mostly expressed on the CD1a-derived DC (40%) as compared to the CD14-derived DC (16%). The CD1a-derived DC have been shown to display features that are associated with Langerhans cells, including the
25 presence of Birbeck granules.

 Next, the in situ distribution of Langerin was examined by immunohistological analysis of various human tissues. Langerin is selectively expressed by LC-like DC. Immunohistological analysis of Langerin expression was
30 made on sections of skin, tonsil and lung. Within the epidermis, Langerin was only found on LC, which also stained with anti-Lag and anti-CD1a antibodies. In skin, staining with mAb DCGM4, or mAb DCGM4 plus anti-Lag or anti-CD1a, showed positive Langerin expression by LC. In
35 tonsil, the Langerin-positive cells were found in the epithelium (e.g., on follicular mantle B cells). A few cells were occasionally stained in the T cell areas, but never observed in germinal centers Langerin⁺ cells were also notably present in lung epithelium. In lung, mAb

TNF- α was determined at various time points, cells were recovered and double-labeled using anti-CD1a-FITC and DCGM4 plus anti-mouse IgG-PE. The results are representative of 5 experiments.

5 No staining was detected at day 0 or day 6, indicating that CD34⁺ HPC and their immediate progeny do not express Langerin. The antigen appeared at day 7, on a small subset of the CD1a⁺ cells. Between day 7 to day 12, Langerin expression reached a maximum, staining
10 between 15 to 35% of CD1a⁺ cells. The dendritic nature of the Langerin-expressing cells in the cultures was confirmed on cytopsin preparations of DCGM4⁺ FACS-sorted cells.

 Caux, et al. (1996) J Exp Med 184:695-706 have
15 further shown that CD34⁺ HPC differentiate along two independent pathways from distinct precursor subsets, identified by mutually exclusive expression of CD1a and CD14 at early time points during the culture (day 5-7). When such precursors were separated by FACS-sorting at
20 day 6 and cultured with GM-CSF and TNF- α for 6 more days, Langerin was found mostly expressed on the CD1a-derived DC (40%) as compared to the CD14-derived DC (16%). The CD1a-derived DC have been shown to display features that are associated with Langerhans cells, including the
25 presence of Birbeck granules.

 Next, the in situ distribution of Langerin was examined by immunohistological analysis of various human tissues. Langerin is selectively expressed by LC-like DC. Immunohistological analysis of Langerin expression was
30 made on sections of skin, tonsil and lung. Within the epidermis, Langerin was only found on LC, which also stained with anti-Lag and anti-CD1a antibodies. In skin, staining with mAb DCGM4, or mAb DCGM4 plus anti-Lag or anti-CD1a, showed positive Langerin expression by LC. In
35 tonsil, the Langerin-positive cells were found in the epithelium (e.g., on follicular mantle B cells). A few cells were occasionally stained in the T cell areas, but never observed in germinal centers. Langerin⁺ cells were also notably present in lung epithelium. In lung, mAb

DCGM4 and counterstaining with hematoxylin showed LC-like Langerin⁺ cells only in bronchiolar epithelium. The Langerin⁺ DC showed dendritic morphology. No staining was detected with control mAbs. These results were representative of 5 experiments.

By contrast to CD34-derived DC, mAb DCGM4 did not react with DC obtained from peripheral blood monocytes cultured 6 days with a combination of GM-CSF and IL-4, thus, further confirming the restriction of Langerin expression. Likewise, Langerin was neither detected in ex-vivo purified DC isolated from peripheral blood, nor in germinal center DC isolated from tonsils.

Finally, mAb DCGM4 was analyzed for reactivity on a panel of different hematopoietic-derived cell types. Langerin was neither detected in ex-vivo isolated T lymphocytes, B lymphocytes, monocytes or granulocytes, nor in myeloid (HL60, KG1, U937, THP1) or lymphoid (Jurkat, JY, PREALP) cell lines.

Taken together, the above data suggest that Langerin expression is restricted to an immature DC compartment, and that it is subsequently lost upon DC maturation.

Example 12

Langerin Expression is Upregulated by TGF- β and Decreased Following CD40 Activation

Since mAb DCGM4 was found to react selectively with LC-like immature DC, it was investigated whether factors that influence DC maturation would affect the expression levels of Langerin.

Studies in vitro and in vivo have shown that TGF- β plays an essential role in LC development. Therefore, the effect of TGF- β on Langerin expression by in vitro derived DC was evaluated.

To do so, cord blood CD34⁺ HPC were cultured for 12 days in GM-CSF and TNF- α in absence or presence of TGF- β from day 7 to day 12. Subsequently, the DC were cultured with L-cells transfected with CD40L for 2 days. Cells were processed for staining without or after pretreatment

09787192-031501

with 0.1% saponin, using mAbs revealed by FITC-conjugated anti-mouse Ig. The results are representative of more than 5 experiments.

- CD34-derived DC were supplemented with TGF- β for the last three days of culture (day 9-12). This resulted in strong upregulation of Langerin expression. In addition to increasing the proportion of Langerin⁺ cells, TGF- β raised the mean number of surface-membrane molecules per cell (93×10^3 instead of 33×10^3 without TGF- β). The effect of TGF- β was predominantly exerted on the CD14-derived DC subset, normally devoid of Langerhans markers such as the Birbeck granule associated antigen Lag. Langerin expression is not restricted to the plasma membrane, but is also detected intracellularly following membrane permeabilization. Levels of intracellular Langerin were also markedly enhanced by TGF- β . It was also found that although TGF- β also increased the expression of the LC markers Lag and E-cadherin, Lag was never detected at the cell-surface.
- Removal of TNF- α , a cytokine known to induce DC maturation, for the last three days of culture, upregulated Langerin expression whether in the presence or absence of TGF- β . In line with this result, a strong decrease in Langerin surface-membrane expression was found associated with an increase of HLA-DR following activation with CD40L, a signal which triggers the maturation of DC including upregulation of costimulatory molecules. Altogether, these results confirm that TGF- β which induces an LC phenotype, upregulates Langerin expression, whereas signals that trigger DC maturation decrease Langerin expression.

Example 13

- Intracellular Langerin and Lag Co-localize in Immature DC, but Dissociate Upon Activation

A unique feature of Langerhans cells is the presence of intracytoplasmic Birbeck granules (BG). As Langerin was found to be selectively expressed in Langerhans-type

DC, its relationship with Lag antigen was further examined. Thus, CD34-derived DC supplemented with TGF- β , and which contain a high proportion of BG⁺ cells as detected by electron microscopy, were analyzed by double fluorescence staining and confocal microscopy. CD34-derived DC were supplemented with TGF- β from day 7 to day 9 of culture. Double-color confocal laser scanning microscopy was performed at day 9 and after two subsequent days of CD40L activation. Langerin-positive and Lag-positive vesicles were found co-localized in day 9 immature DC. After CD40L activation, Langerin⁺Lag⁻ staining was observed near the nucleus. No co-localization was detected between Langerin and HLA-DR or LAMP-1, irrespective of CD40L activation. Langerin and Lag were found to display a remarkable intracellular co-localization. Strikingly however, localization of the two markers segregated upon activation with CD40L. Thus, Langerin was found in the absence of Lag in close proximity to the nucleus, whereas a co-localized expression of the two markers only remained immediately beneath the surface membrane.

These results indicate that Langerin is distinct from Lag, and that Langerin is routed towards a deep cellular compartment during DC activation. Finally, Langerin was never found to co-localize with the lysosomal marker LAMP-1, nor with HLA-DR, irrespective of activation by CD40 crosslinking.

Example 14

30 Langerin is Associated with Endocytic Structures and Birbeck Granules

Since Langerin is expressed at the cell surface (as detected by FACS in the absence of membrane permeabilization) electron microscopy was performed on epidermal cell suspensions to analyze its precise distribution. Langerhans cells are easily recognizable in such suspensions, by their folded nuclei, lack of keratin

filaments, lack of desmosomes and melanosomes, and the presence of characteristic BG.

An epidermal cell suspension was obtained as described above. mAbs were revealed by 5 nm gold-labeled goat anti-mouse IgG1. CD1a is homogeneously distributed at the cell surface, whereas Langerin is often associated with areas of membrane thickening. Cytoplasmic sandwiching structures and coated pits were visualized upon staining with DCGM4 at 4°C. Upon incubation at 37°C, cytoplasmic gold particle-containing coated vesicles and Birbeck granules were seen. DCGM4 staining and gold particles were revealed on the luminal side of BG. These results were representative of 3 experiments on different skin samples.

Staining at 4°C with DCGM4 and 5 nm gold particles confirmed that Langerin is clearly associated with the cell surface in LC, although at a lower density than CD1a (161.5 ± 97.1 versus 1589.1 ± 418.8 gold granules/100µm membrane). In addition to single gold particles, a spontaneous clustering was observed with DCGM4, even though the antibody was ultracentrifuged. An isotype-matched control mouse IgG1 did not bind to the LC (1.3 ± 3.4 granules/100 µm). No labeling of DCGM4 was observed on the keratinocytes or melanocytes present in the epidermal cell suspensions.

As compared to the rather homogeneous distribution of CD1a, Langerin was not randomly distributed at the cell surface, but was often associated with particular areas of membrane thickening. Furthermore, DCGM4 induced typical endocytic coated pits at the cell membrane at 4°C. The number of coated pits was significantly enhanced (an average 3.6 times) by DCGM4, as compared to staining with anti-CD1a or control mouse IgG1. Notably, the coated pits induced during DCGM4 staining contained gold particles. Furthermore, when cells were allowed to warm up before fixation, DCGM4 staining was observed inside coated vesicles already after 2 min. at 37° C. These data demonstrate that Langerin is associated with

09787197-031504

structures characteristic of early steps of receptor-mediated endocytosis.

Staining with DCGM4 at 4° C also resulted in the formation of cytomembrane sandwiching structures at the cell surface. Consistently, following DCGM4 at 4° C, gold labeled BG were seen in continuity with the cell membrane, and found inside the cytoplasm when cells were warmed up to 37° C for 2 min. The BG were labeled in their central striated lamella or in their bulb.

Taken together, these results indicate that Langerin is associated with endocytic structures and can also gain access to Birbeck granules from the cell membrane.

Example 15

Langerin Mediates Rapid Internalization in DC

To further examine the role of Langerin in endocytosis, we analyzed its capacity to internalize DCGM4 as a ligand.

CD34-derived DC supplemented with TGF- β were labeled at 4° C with mAbs and subsequent F(ab')₂ biotinylated secondary antibody. Cells were incubated at 37° C for time periods indicated, and internalization measured as decreased cell-surface-bound antibody determined by FACS analysis using PE-conjugated streptavidin. MAb DCGM4 is rapidly internalized at 37°C, with similar kinetics as an anti-mannose-receptor mAb (positive control). In fixed cells, no decrease of cell-surface fluorescence was detected. Results were analyzed as the percentage decrease of mean fluorescence intensity (MFI), as compared to control samples kept at 4° C.

It was found that the antibody was very rapidly internalized by DC at 37° C, but not at 4° C. Approximately 75% of surface-membrane bound DCGM4 was internalized already within one minute at 37°C, with similar kinetics to that of anti-mannose-receptor mAb DCGM1, used as positive control for receptor-mediated endocytosis. In a representative experiment, the half-life ($t_{1/2}$) of DCGM4 at the cytomembrane was calculated

to be 4.5 minutes at 37° C, with an internalization rate of $k=15.3\%/min$. The rapid disappearance of mAb DCGM4 from the cell-surface was not due to antibody dissociation, as no decrease in fluorescence was observed in glutaraldehyde-fixed DC incubated at 37° C. Finally, Langerin did not display a mannose-type receptor specificity, as DCGM4 failed to inhibit uptake of Dextran-FITC at 37° C and binding of the antibody was not inhibited by mannan at 4° C.

These results are in accordance with the above electron microscopy analysis and demonstrate that Langerin is implicated in a rapid endocytosis process by DC.

Example 16

Triggering of Cell-Surface Langerin Results in Birbeck Granule Formation

As triggering of cell surface Langerin resulted in the formation of CMS and the detection of gold-labeled BG as visualized by electron microscopy, the potential role of Langerin cell surface in BG formation was further investigated. An epidermal cell suspension was obtained as described above. Cells were incubated with an excess of mAb DCGM4 or anti-CD1a at 4°C, and processed immediately for electron microscopy, or left at room temperature for 5 min. before fixation. LC incubated with DCGM4 displayed a striking accumulation of Birbeck granules in the perinuclear region. The effect is predominantly observed at 4°C, as subsequent warming up results in vacuolization. Treatment with anti-CD1a mAb failed to induce significant changes in the LC cytoplasm, whether in cells kept at 4°C or brought up to room temperature.

This treatment resulted in a considerable increase of densely packed BG in the LC cytoplasm, with a marked accumulation in the perinuclear region around the Golgi. The BG displayed an elongated, round, or irregularly-shaped expanded portion, in addition to their typical

rod-shaped part. Moreover, the cytoplasm of DCGM4-treated LC was filled with numerous rounded or elongated vesicles.

When LC were warmed up to room temperature (5 min.) following incubation with excess DCGM4, increased fusion was observed between single and short rod-shaped BG and large vesicular components. Moreover, numerous vesicles of various sizes and shapes occupied a considerable volume of the LC cytoplasm.

In contrast to DCGM4, incubation of epidermal cell suspensions with an excess anti-CD1a mAb only led to the formation of some small vesicles but no other significant changes in the LC cytoplasm. Notably, no accumulation of perinuclear BG was observed, even when cells were allowed to warm up before fixation. Similarly, incubation with a control mouse IgG1 or with anti-E-cadherin mAb did not modify the BG granules.

Taken together, these data demonstrate that cell-surface Langerin actively participates in BG formation, resulting in their perinuclear accumulation.

Example 17

Langerin is a 40 kDa N-glycosylated Protein

Immunoprecipitation with DCGM4 from DC extracts and, subsequent elution with SDS-PAGE sample buffer yielded a homogeneous band of 40-42 kDa molecular mass. SDS-PAGE analysis of immunopurified Langerin in non-reducing and reducing conditions was carried out. If DTT was omitted all along the purification steps, the profile was not modified on the gel, suggesting that Langerin is present at the cell membrane as a single chain or as an homodimer with non covalent association.

2-D analysis of immunopurified Langerin was conducted establishing the molecular mass of the molecule and indicating that Langerin had a pI of 5.2-5.5. Finally, a dot-blot analysis of Langerin using (1) creatinase from E. coli as non-glycosylated control, (2) transferrin as positive N-glycosylated control, (3) N-

glycosidase, (4) Langerin, and (5) N-glycosidase-treated Langerin, demonstrated that Langerin is a glycoprotein, and that most of the carbohydrate constituents were removed by N-glycosylase treatment.

5

Example 18

Isolating a Nucleic Acid Encoding a Langerin

10 Numerous methods are available to isolate a gene encoding a purified protein, especially where antibodies which recognize the protein exist. One method is to determine methods for purification of the protein and subsequently to determine the peptide sequences. Given sufficient sequence information, and using redundant
15 oligonucleotides, PCR or hybridization techniques will allow for isolation of genes encoding Langerin proteins.

Another alternative is to generate additional antibodies to Langerin proteins, which may be isolated by immunoaffinity methods using the DCGM4 antibodies. See
20 above. These antibodies are applicable in "panning" techniques, such as described by Seed and Aruffo (1987) Proc. Nat'l Acad. Sci. USA 84:3365-3369. Phage expression techniques are also applicable to screen cDNA libraries derived from appropriate DC or T cell
25 subpopulations enriched for Langerin expression. Glycosylation interference with antibody recognition will be generally less problematic in the phage selection systems. Cell sorting techniques on a mammalian expression library are applicable also.

30 Another method for screening an expression library is to use antibody to screen successive subpopulations of libraries. The following provides one method of screening using small populations of cells on slides stained by a specific labeling composition, e.g., an
35 antibody.

For example, on day 0, precoat 2-chamber permanox slides with 1 ml per chamber of fibronectin, 10 ng/ml in PBS, for 30 min. at room temperature. Rinse once with

09707192-031501

PBS. Then plate COS cells at $2-3 \times 10^5$ cells per chamber in 1.5 ml of growth media. Incubate overnight at 37°C .

On day 1 for each sample, prepare 0.5 ml of a solution of 66 $\mu\text{g/ml}$ DEAE-dextran, 66 μM chloroquine, and 4 μg DNA in serum free DME. For each set, a positive control is prepared, e.g., of human IL-10-FLAG cDNA construct at 1 and 1/200 dilution, and a negative mock. Rinse cells with serum free DME. Add the DNA solution and incubate 5 h at 37°C . Remove the medium and add 0.5 ml 10% DMSO in DME for 2.5 min. Remove and wash once with DME. Add 1.5 ml growth medium and incubate overnight.

On day 2, change the medium. On days 3 or 4, the cells are fixed and stained. Rinse the cells twice with Hank's Buffered Saline Solution (HBSS) and fix in 4% paraformaldehyde (PFA)/glucose for 5 min. Wash 3X with HBSS. The slides may be stored at -80°C after all liquid is removed. For each chamber, 0.5 ml incubations are performed as follows. Add HBSS/saponin (0.1%) with 32 $\mu\text{l/ml}$ of 1M NaN_3 for 20 min. Cells are then washed with HBSS/saponin 1X. Soluble antibody, e.g., DCGM4, is added to cells and incubate for 30 min. Wash cells twice with HBSS/saponin. Add second antibody, e.g., Vector anti-mouse antibody, at 1/200 dilution, and incubate for 30 min. Prepare ELISA solution, e.g., Vector Elite ABC horseradish peroxidase solution, and preincubate for 30 min. Use, e.g., 1 drop of solution A (avidin) and 1 drop solution B (biotin) per 2.5 ml HBSS/saponin. Wash cells twice with HBSS/saponin. Add ABC HRP solution and incubate for 30 min. Wash cells twice with HBSS, second wash for 2 min., which closes cells. Then add Vector diaminobenzoic acid (DAB) for 5 to 10 min. Use 2 drops of buffer plus 4 drops DAB plus 2 drops of H_2O_2 per 5 ml of glass distilled water. Carefully remove chamber and rinse slide in water. Air dry for a few minutes, then add 1 drop of Crystal Mount and a cover slip. Bake for 5 min. at $85-90^\circ \text{C}$.

Alternatively, the Langerin proteins are used to affinity purify or sort out cells expressing the ligand.

09787197-0315001
105130 26728760

See, e.g., Sambrook, et al. (supra) or Ausubel, et al., (supra) both of which are incorporated herein by reference.

5

Example 19

Human Genomic Langerin

The genomic sequence (SEQ ID NO: 3) and the exon-intron organization of human Langerin has now been
 10 determined. The genomic sequence includes the Langerin cDNA sequence (SEQ ID NO: 1) as 6 exons separated by 5 introns. The nucleotide sequences of exons 1-6 are provided in SEQ ID NOS 4-9 respectively. This 10,663 nucleotide genomic sequence includes 3251 and 1808
 15 nucleotides at the 5' and 3' ends of exons 1 and 6 respectively, and is predicted to include the sequence of the human Langerin promoter.

Exon 1 (SEQ ID NO: 4) corresponds to positions 3252-3371 of SEQ ID NO: 3. Exon 2 (SEQ ID NO: 5) corresponds
 20 to positions 3467-3583 of SEQ ID NO: 3. Exon 3 (SEQ ID NO: 6) corresponds to positions 5051-5425 of SEQ ID NO: 3. Exon 4 (SEQ ID NO: 7) corresponds to positions 6020-6171 of SEQ ID NO: 3. Exon 5 (SEQ ID NO: 8) corresponds to positions 7252-7370 of SEQ ID NO: 3. Exon 6 (SEQ ID
 25 NO: 9) corresponds to positions 7871-8855 of SEQ ID NO: 3.

Human Langerin is a 328 amino acid transmembrane type II Ca++ dependent lectin. Amino-acid translation of the 6 exons shows that exon 1 encodes the NH2-terminal
 30 portion of the intracellular domain (24 amino acids), while both the membrane-proximal portion of the intracellular domain (19 amino acids) and the entire transmembrane domain (20 amino acids) are encoded by exon 2. Exon 3 encodes the entire membrane-proximal non-
 35 carbohydrate recognition portion of the extracellular domain (125 amino acids). The extracellular carbohydrate-recognition domain (CRD) is encoded jointly by exons 4 (51 amino acids), 5 (39 amino acids), and the beginning of exon 6 (50 amino acids) the latter terminating the
 40 COOH-portion of the molecule. Notably, the CRD of

09787192-031504

Langerin shares the three exon structure of other type II transmembrane lectins including the Kupffer cell receptors, rat hepatic lectin, and the CD23 low-affinity IgE receptor, with the coding sequences interrupted by
5 introns at analogous positions.

The availability of the genomic Langerin sequence now allows for the identification of human Langerin promoter elements, expected to be situated within the 500-600 nucleotides 5' of the ATG initiation codon at
10 position 48 of exon 1. By analogy, in the rat Kupffer cell receptor, homologous to human Langerin, the transcription initiation site is situated 51 bases upstream from the ATG translation initiation codon. Two nucleotide sequences have been identified 5' to the start
15 of transcription, that are candidates for regulation of gene expression. One is a 54-base tandemly repeated sequence from nucleotides -151 to -98 and -97 to -44, and the other is a heptanucleotide sequence (GAGGCAG) that is repeated 4 times in 380 bases 5' to the start site (G.W.
20 Hoyle and R.L. Hill, J. Biol Chem, 1991, 266:1850-1857).

Human Langerin is specifically expressed in Langerhans cells, a subtype of immature dendritic cells specialized in capture and processing of antigen. Thus, it will be feasible to construct DNA sequences encoding
25 antigens of choice for targeting selective expression in Langerhans cells under control of the Langerin promoter elements.

Human Langerin sequence enables the identification of the mouse homolog Langerin genomic sequence. This
30 information will be key to the construction of mice deficient in the Langerin gene, to further explore the function of Langerin and of the Birbeck granules, the unique endocytic organelles of Langerhans cells that are induced by engagement of Langerin.

35 The human Langerin sequence also enables the sequencing of possible mutations in the Langerin gene in patients suffering from immunological disorders of unknown etiology.

097827487-001504

Example 20

Chromosomal Mapping of Human Langerin

Chromosomal localization was performed by radiation
5 hybrid (RH) mapping (D.R. Cox et al., Science, 1990, 250:
245-250), using the Stanford G3 Radiation Hybrid panel
(Research Genetics, Inc., Huntsville, AL). Briefly, two
oligonucleotides (U863 and L1180) used to amplify
10 Langerin cDNA by PCR reaction were selected as they also
amplified a stretch of human genomic DNA. PCR reactions
were carried out with U863 / L1180 on the Stanford G3
panel of 83 clones covering the human genome. PCR data
were submitted to the Stanford Human Genome Center
RHserver (LIENHYPERTEXTE
15 mailto:rhserver@paxil.stanford.edu
rhserver@paxil.stanford.edu) for mapping using the RHMAP
statistical program (RHMAP).

Results from the RH Server indicated as closest
matches the markers SHGC-58922 (LOD score: 8.67) and
20 SHGC-12714 (LOD score 7.74). Both markers are located on
chromosome 2, linked to the GDB locus D2S292 (SHGC source
AFH203yb6t: SHGC-1610 microsatellite marker). Genes
mapping near the D2S292 locus include: transforming
growth factor-alpha precursor (2p13), dynactin (2p13),
25 gamma actin enteric smooth muscle form (2p13), annexin IV
(2p13), MAD (2p12-13), alpha-CPI, early growth response
4, nucleolysin TIA-1, protein tyrosine phosphatase P,
protein kinase C substrate 80K-H, glutamine-fructose 6-
phosphate transaminase, retinoic acid-responsive protein,
30 RAB.1A, and pleckstrin. The results localize the gene of
human Langerin to chromosome 2p13, in the vicinity of the
D2S292 locus.

The localization of human Langerin will permit to
survey immunological genetic disorders for a potential
35 association with the chromosome 2p13 region.

00737192-031501

Example 21

Identification of Mouse Langerin

- Using the nucleotide sequence of the human Langerin, a search was performed in the NCBI database. This permitted to identify two mouse ESTs (AA764540 and AA423304) displaying considerable homology to the human sequence. These ESTs were then used to extend the sequence, using RACE-PCR on a mouse lung cDNA library: The entire coding region, with the exception of the extremity of the 5' end was determined. Nine nucleotides completing the sequence were subsequently identified by sequencing a mouse cosmid clone. A contig of 1756 nucleotides was obtained and is shown in SEQ ID NO: 10.
- An open-reading frame of 978 nucleotides was identified (positions 266-1243 in the contig of SEQ ID NO: 10), encoding a predicted transmembrane type II Ca++ dependent lectin of 326 amino acids (SEQ ID NO: 11). The mouse amino acid sequence presents considerable homology (66.6%) with human Langerin (alignement shown in Table 1), with conservation of the key structural features of the molecule (ie. intracytoplasmic proline-rich motif as potential signal transduction site, and extracellular carbohydrate-recognition domain (CRD) with an EPN motif as found in human Langerin and indicative of mannose-binding specificity). These data indicate that the above identified mouse molecule is the homolog of human Langerin.
- The mouse Langerin protein has been successfully expressed in mammalian cells (murine COP5 fibroblasts), which have been used for immunization of mice to produce monoclonal antibodies (mAbs). A number of mAbs recognizing mouse Langerin protein have been isolated 7 of these mAbs display crossreactivity on human Langerin. The epitopes recognized by the cross-reactive mAbs have all been mapped to the intracytoplasmic domain using truncated forms of recombinant human Langerin protein. This is consistent with the remarkably high conservation

Table 1

		MLV E E E A P D A H F T V D K Q N I S L W P R E P P P K S G L S L V L G K T L	Majority
		10 20 30 40	
1	MTVEKEAPDAHFTVDKQNI SLWPREP PPKSGPSLVFGKTP	hu Langerin aa	
1	ML--EEAPEAHFTVDKQNI SLWPREP PPK QDLSFVLRKPL	mouse langerin aa	
		TVRAALICLALVLVASVVLQAVLYPRLMGTIL DVKSDAQL	Majority
		50 60 70 80	
41	TVRAALICLTLVLVASVVLQAVLYP R F M G T I S D V K T N V Q L	hu Langerin aa	
39	CICVAFTCLALVLVTSIVLQAVFYPR LMGKIL DVKSDAQ M	mouse langerin aa	
		LKGRVDNISTLGSDLKTESGGVDAAGVQIQIVNTSLGRVR	Majority
		90 100 110 120	
81	LKGRVDNISTLDSEIKXNSDGM E A A G V Q I Q M V N E S L G Y V R	hu Langerin aa	
79	LKGRVDNISTL GSDLKTERGRVDDAEVQM Q I V N T T L K R V R	mouse langerin aa	
		SQILSLET SVEIANAQLLILTRSWGEVSSLSAQIPELKSD	Majority
		130 140 150 160	
121	SQFLKLLKTSVEKANAQIQILTRSWEEVSTLNAQIPELKSD	hu Langerin aa	
119	SQILSLETSMKIAN D Q L L I L T M S W G E V D S L S A X I P E L K R D	mouse langerin aa	
		LDKASALNTKVQGLQGSLENVSKLLKQQSDILEVVAQGWK	Majority
		170 180 190 200	
161	LKASALNTKIRALQGSLENMSKLLK R Q N D I L Q V V S Q G W K	hu Langerin aa	
159	LDKASALNTK VQGLQNSLENVNKL L K Q Q S D I L E M V A R G W K	mouse langerin aa	
		YFSGNFYFPSLIPKTNWYSAEQFCVSRNAHLTSVSSSESEQE	Majority
		210 220 230 240	
201	YFKGNFYFPSLIPKTNWYSAEQFCVSRNSHLTSVTSESEQE	hu Langerin aa	
199	YFSGNFYFPS R T P K T N W Y S A E Q F C I S R K A H L T S V S S E S E Q K	mouse langerin aa	
		FLYKAAAGGLIHWIGLTKAGSEGDWSWVDDTSFNKVQSARF	Majority
		250 260 270 280	
241	FLYKTAGGLIYWIGLTKAGMEGDWSWVDDTFPNKVQSARF	hu Langerin aa	
239	FLYKAAADGIPHWIGLTKAGSEGDWYVVDQTSFNKEQSRRF	mouse langerin aa	
		WIPGEPNNAAGNNEHCGNIKASALQAWNDGPGCDNTFLFICK	Majority
		290 300 310 320	
281	WIPGEPNNAAGNNEHCGNIKAPSLQAWNDAPCDKTFLFICK	hu Langerin aa	
279	WIPGEPNNAAGNNEHCANIRVSALKCWNDGPGCDNTFLFICK	mouse langerin aa	
		RPYVQST E	Majority
321	RPYVVPSEP	hu Langerin aa	
119	RPYVQTTE	mouse langerin aa	

09787192-031501

in the cytoplasmic tail (first 30 residues) of human and mouse Langerin (see Table 1).

Availability of the sequence encoding mouse Langerin will permit the performance of studies (RT-PCR, Northern, in situ hybridization) to determine the expression pattern of Langerin in various tissues. Furthermore, the current sequence can be extended to identify the genomic DNA in order to construct mice deficient in the Langerin gene. Such animals should be highly valuable to further understand the role of Langerin in the regulation of the immune response. MABs directed against mouse Langerin will be useful to study expression of the protein in mouse tissues. Also, mAb can be injected in mice to analyze the impact of triggering Langerin on the dendritic cell system in vivo. These studies will be important to complement efforts to target antigen (ie. tumor antigens) to dendritic cells in vivo via their cell-surface Langerin with the aim of inducing protective antitumor immunity. Targeting experiments can, for instance, be performed by injecting mice (prior of after challenge with tumor) with mAb against mouse Langerin to which tumor antigen has been coupled chemically.

Many modifications and variations of this invention can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. The specific embodiments described herein are offered by way of example only, and the invention is to be limited only by the terms of the appended claims, along with the full scope of equivalents to which such claims are entitled.

WHAT IS CLAIMED IS:

1. An isolated antibody which selectively binds a
5 mammalian polypeptide, said polypeptide characterized
by:
- a) binds with selectively to DCGM4 monoclonal
antibody;
 - b) natural form of said polypeptide has a molecular
10 weight by SDS PAGE of ~40 kD; and
 - c) expressed on the cell surface of Langerhans
cells.
2. The antibody of claim 1 which is a monoclonal
15 antibody.
3. A hybridoma that produces a monoclonal antibody of
claim 2.
- 20 4. The hybridoma of claim 3, which has ATCC Accession
No. HB-12576.
5. A method for analyzing a cell population, comprising
detecting or measuring the presence of Langerin with said
25 antibody of claim 1.
6. A substantially pure mammalian protein characterized
by:
- a) a molecular weight by SDS PAGE of approximately
30 40 kDa with natural glycosylation; and
 - b) binds selectively to said antibody of claim 1.
7. A protein having an amino acid sequence shown in SEQ
ID NO: 2.
- 35 8. A protein having an amino acid sequence shown in SEQ
ID NO: 11.
9. A fragment of the protein of claim 7, which:

09787192-031501

- a) expresses an immunological epitope of said Langerin; or
- b) modulates an immune response.

5 10. A nucleic acid sequence which encodes a protein shown in SEQ ID NO: 2.

11. The nucleic acid sequence of claim 10 comprising the coding region shown in SEQ ID NO: 1.

10

12. A nucleic acid sequence which encodes a protein shown in SEQ ID NO: 11.

13. The nucleic acid sequence of claim 12 comprising the
15 coding region shown in SEQ ID NO: 10.

14. A nucleic acid sequence shown in SEQ ID NO: 3 or a fragment thereof.

20 15. A nucleic acid sequence comprising any one of SEQ ID NOS: 4-9.

105710-26128200

ANTIBODIES TO MAMMALIAN DC ANTIGENS AND USES

5

ABSTRACT

Purified mammalian DC cell surface protein,
designated Langerin, nucleic acids encoding Langerin, and
antibodies which specifically bind Langerin.

10

09787192-034501

SEQUENCE LISTING

<110> Schering Corporation

<120> Antibodies to Mammalian DC Antigen and Uses

<130> SF0820K1

<140> EP 99 400 394.5

<141> 1999-02-18

<150> EP 98 402 374.7

<151> 1998-09-25

<160> 11

<210> 1

<211> 1547

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (56) ... (1039)

<400> 1

ctggaaaggg cagccagaag cacctgtgct cccaggataa ggggtgagcac tcagg atg 58
Met
1

act gtg gag aag gag gcc cct gat gcg cac ttc act gtg gac aaa cag 106
Thr Val Glu Lys Glu Ala Pro Asp Ala His Phe Thr Val Asp Lys Gln
5 10 15

aac atc tcc ctc tgg ccc cga gag cct cct ccc aag tcc ggt cca tct 154
Asn Ile Ser Leu Trp Pro Arg Glu Pro Pro Pro Lys Ser Gly Pro Ser
20 25 30

ctg gtc cgg ggg aaa aca ccc aca gtc cgt gct gca tta atc tgc ctg 202
Leu Val Pro Gly Lys Thr Pro Thr Val Arg Ala Ala Leu Ile Cys Leu
35 40 45

acg ctg gtc ctg gtc gcc tcc gtc ctg ctg cag gcc gtc ctt tat ccc 250
Thr Leu Val Leu Val Ala Ser Val Leu Leu Gln Ala Val Leu Tyr Pro
50 55 60 65

cgg ttt atg ggc acc ata tca gat gta aag acc aat gtc cag ttg ctg 298
Arg Phe Met Gly Thr Ile Ser Asp Val Lys Thr Asn Val Gln Leu Leu
70 75 80

aaa ggt cgt gtg gac aac atc agc acc ctg gat tct gaa att aaa aag 346
Lys Gly Arg Val Asp Asn Ile Ser Thr Leu Asp Ser Glu Ile Lys Lys
85 90 95

09787192-031501

aat agt gac ggc atg gag gca gct ggc gtt cag atc cag atg gtg aat Asn Ser Asp Gly Met Glu Ala Ala Gly Val Gln Ile Gln Met Val Asn 100 105 110	394
gag agc ctg ggt tat gtg cgt tct cag ttc ctg aag tta aaa acc agt Glu Ser Leu Gly Tyr Val Arg Ser Gln Phe Leu Lys Leu Lys Thr Ser 115 120 125	442
gtg gag aag gcc aac gca cag atc cag atc tta aca aga agt tgg gaa Val Glu Lys Ala Asn Ala Gln Ile Gln Ile Leu Thr Arg Ser Trp Glu 130 135 140 145	490
gaa gtc agt acc tta aat gcc caa atc cca gag tta aaa agt gat ttg Glu Val Ser Thr Leu Asn Ala Gln Ile Pro Glu Leu Lys Ser Asp Leu 150 155 160	538
gag aaa gcc agt gct tta aat aca aag atc cgg gca ctc cag ggc agc Glu Lys Ala Ser Ala Leu Asn Thr Lys Ile Arg Ala Leu Gln Gly Ser 165 170 175	586
ttg gag aat atg agc aag ttg ctc aaa cga caa aat gat att cta cag Leu Glu Asn Met Ser Lys Leu Lys Arg Gln Asn Asp Ile Leu Gln 180 185 190	634
gtg gtt tct caa ggc tgg aag tac ttc aag ggg aac ttc tat tac ttt Val Val Ser Gln Gly Trp Lys Tyr Phe Lys Gly Asn Phe Tyr Tyr Phe 195 200 205	682
tct ctc att cca aag acc tgg tat agt gcc gag cag ttc tgt gtg tcc Ser Leu Ile Pro Lys Thr Trp Tyr Ser Ala Glu Gln Phe Cys Val Ser 210 215 220 225	730
agg aat tca cac ctg acc tcg gtg acc tca gag agt gag cag gag ttt Arg Asn Ser His Leu Thr Ser Val Thr Ser Glu Ser Glu Gln Glu Phe 230 235 240	778
ctg tat aaa aca gcg ggg gga ctc atc tac tgg att ggc ctg act aaa Leu Tyr Lys Thr Ala Gly Gly Leu Ile Tyr Trp Ile Gly Leu Thr Lys 245 250 255	826
gca ggg atg gaa ggg gac tgg tcc tgg gtg gat gac acg cca ttc aac Ala Gly Met Glu Gly Asp Trp Ser Trp Val Asp Asp Thr Pro Phe Asn 260 265 270	874
aag gtc caa agt gcg agg ttc tgg att cca ggt gag ccc aac aat gct Lys Val Gln Ser Ala Arg Phe Trp Ile Pro Gly Glu Pro Asn Asn Ala 275 280 285	922
ggg aac aat gaa cac tgt ggc aat ata aag gct ccc tca ctt cag gcc Gly Asn Asn Glu His Cys Gly Asn Ile Lys Ala Pro Ser Leu Gln Ala 290 295 300 305	970
tgg aat gat gcc cca tgt gac aaa acg ttt ctt ttc att tgt aag cga Trp Asn Asp Ala Pro Cys Asp Lys Thr Phe Leu Phe Ile Cys Lys Arg 310 315 320	1018
ccc tat gtc cca tca gaa ccg tga c aggacaggct cccaagctca Pro Tyr Val Pro Ser Glu Pro 325	1063
ctctttgagc tccaacgctt gttaaacatg aggaatgcc tctttcttcc ccgaactcca	1123

09787192.031504

ggatgacttt gcacgttaat ttttcttgct tcaaaattgt cccacagtgg cattctggag 1183
 tccgtctgtc ttggctggaa attctctgac gtcttggagg cagctggaat ggaaaggaga 1243
 attcaggta aagtgaggagg ggtgggtaga gaggatttag aagttccaat tgccttgcta 1303
 aggaggatca agaccctgaa tccggcacaa caccctgggg ttttccactc tttcagagaa 1363
 acctcagctt catcacatca aagttactcc agagcaacca agcaattctc ctgatatgtt 1423
 catccagggc ttttcttggc caaaccctc agaatttcca tgtctctgct tagctgtgct 1483
 ggcagctagc agctggctgt gtttgcagtg caaatagctc tgttcttgga aatcctgctc 1543
 atgg 1547

<210> 2
 <211> 328
 <212> PRT
 <213> Homo sapiens

<400> 2
 Met Thr Val Glu Lys Glu Ala Pro Asp Ala His Phe Thr Val Asp Lys
 1 5 10 15
 Gln Asn Ile Ser Leu Trp Pro Arg Glu Pro Pro Pro Lys Ser Gly Pro
 20 25 30
 Ser Leu Val Pro Gly Lys Thr Pro Thr Val Arg Ala Ala Leu Ile Cys
 35 40 45
 Leu Thr Leu Val Leu Val Ala Ser Val Leu Leu Gln Ala Val Leu Tyr
 50 55 60
 Pro Arg Phe Met Gly Thr Ile Ser Asp Val Lys Thr Asn Val Gln Leu
 65 70 75 80
 Leu Lys Gly Arg Val Asp Asn Ile Ser Thr Leu Asp Ser Glu Ile Lys
 85 90 95
 Lys Asn Ser Asp Gly Met Glu Ala Ala Gly Val Gln Ile Gln Met Val
 100 105 110
 Asn Glu Ser Leu Gly Tyr Val Arg Ser Gln Phe Leu Lys Leu Lys Thr
 115 120 125
 Ser Val Glu Lys Ala Asn Ala Gln Ile Gln Ile Leu Thr Arg Ser Trp
 130 135 140
 Glu Glu Val Ser Thr Leu Asn Ala Gln Ile Pro Glu Leu Lys Ser Asp
 145 150 155 160
 Leu Glu Lys Ala Ser Ala Leu Asn Thr Lys Ile Arg Ala Leu Gln Gly
 165 170 175
 Ser Leu Glu Asn Met Ser Lys Leu Leu Lys Arg Gln Asn Asp Ile Leu
 180 185 190

09787192-034501

Gln Val Val Ser Gln Gly Trp Lys Tyr Phe Lys Gly Asn Phe Tyr Tyr
195 200 205

Phe Ser Leu Ile Pro Lys Thr Trp Tyr Ser Ala Glu Gln Phe Cys Val
210 215 220

Ser Arg Asn Ser His Leu Thr Ser Val Thr Ser Glu Ser Glu Gln Glu
225 230 235 240

Phe Leu Tyr Lys Thr Ala Gly Gly Leu Ile Tyr Trp Ile Gly Leu Thr
245 250 255

Lys Ala Gly Met Glu Gly Asp Trp Ser Trp Val Asp Asp Thr Pro Phe
260 265 270

Asn Lys Val Gln Ser Ala Arg Phe Trp Ile Pro Gly Glu Pro Asn Asn
275 280 285

Ala Gly Asn Asn Glu His Cys Gly Asn Ile Lys Ala Pro Ser Leu Gln
290 295 300

Ala Trp Asn Asp Ala Pro Cys Asp Lys Thr Phe Leu Phe Ile Cys Lys
305 310 315 320

Arg Pro Tyr Val Pro Ser Glu Pro
325

<210> 3

<211> 10663

<212> DNA

<213> homo sapiens

<400> 3

tcctgtgtccc	cactccgcgaa	tttgggtgttc	actctactttt	cctccatgac	ttgactgagg	60
gacttgggtg	actctccctc	ggagatgact	cttggcactg	ctggaggcatt	tcatacaacc	120
caggtctctg	ttccagttct	gttgattaaa	cccatgggttc	tcctccaagc	ataaggctgg	180
gcccaaccat	agtctcaacc	agagaaaagg	aaaaagtgc	caagctgttt	tcctcctcca	240
tcctttatcca	gagaaaaaat	tagaacccca	ggaacacagg	aagaggagaa	actaaaaaaa	300
tgagcacaga	cttcccgttc	tcatacacag	gggtattttt	ttggctcacta	ggctctggcca	360
gagttctctc	tctgttttga	agagcagggc	attgtctgag	tgcccgggag	aactgcccac	420
cttctcgact	ggaaaagtgtg	gccaaaggcac	ctgcctgctt	ctctgtcttt	tctcctgccca	480
ggccaggtag	cagatagagc	atccagaggt	ttgcacagga	gaggtgtctca	ggaaaagacct	540
gtagatgagc	atgataaatg	gaagatggct	tctgactccc	tcctctctcca	caaagtggga	600
aatgaaactt	cctatgtgtt	gggactttta	catccacctc	ctctgaaaac	ccagaaggcc	660
cacagcccac	ggattcccatg	cttttggctc	agcttctcca	tgctggaaca	gcctttccgc	720
tagcggtagc	tattctaacct	acctcagcct	gcatttccat	cacttggagt	aaatgctccc	780
tgctccagcc	ccttctcatg	ccatggggcca	gcgcgtaagg	cttctctcatc	ctttcctggg	840
accaatgaga	gggcattctgg	ggggctcttat	aaagtataac	tcataccaca	aaccagcaga	900
gaagaaaccaa	cttctcgacc	gagcgggaaat	ggacaagaca	cttcccaggc	acactgtgtg	960
gcggctctctt	tcctcgcaaat	gcctcaggag	gagacatgga	ccctcgcaag	gccctctcta	1020
ccagggtctt	gagttaattg	gattcttctc	tatccccagg	cccagaaata	ggacaggcta	1080
tgatgaaagg	gccaggagcc	aggagcagaa	gggcagagat	gattctggag	ttaggctggg	1140
ttccaagcat	gagagtcacg	aagtggaaaca	ttctaactct	gtctaagccc	tgatggaaac	1200
aaggctctgc	aaccacacga	gccacctgaa	gagagtagga	attctcatgg	acacccagag	1260
gcaggcagtg	atgtggagcc	aggccctcat	attcggggact	atcaaatgtt	aaaattttag	1320
gcagacagca	ggcctggaaag	ttacagacag	gctgggttagg	gaagcgatgg	cacagaacaa	1380
atgagacacc	atctttgcta	cccagtttcc	ttcccagagc	ttcccagagt	ggggctctct	1440
ctgggtgtccc	tcactccagtc	cgagaagcca	ctccagcctg	aggcccgggt	agctggagag	1500
agagcatcaa	gggcctgttaa	ctacaatgag	ctgcacatag	gcattgggctg	agcgtgcagc	1560

cactatcctc	ccaagtacca	aaggtggcag	tagccctcta	aagcataggt	agatatcttc	1620
ttggccctgt	taacttaacc	tctaatcctg	gtccctgata	caggctgtca	tcacagctcca	1680
ctggctccagc	ctccagtcagc	ccaggacagc	ttctcttttc	ttggccccaga	gcttttgttc	1740
attctctgggt	aaggagtaaa	tcacactctc	ccttatgtct	ttgcattgaa	gatgaatgag	1800
aactctctgga	gacagtgga	aaaagcagca	gctctagaa	ggcagatccc	agggcccaga	1860
ctgtgataat	gtctgagtgg	catgatcagc	tatcaagctc	ccaagtcact	taccattcac	1920
acatccatcc	atgcatactc	tatccatcca	ttcatccatt	caccatcctc	ccaattatct	1980
ggaccaattt	ccttcttttt	tcttctcttc	cctccctctc	tctctctctc	ttctctatat	2040
tccaagaaat	atttgtctat	agtcgtccat	gtgccaattt	tctcagattt	ggcagagctc	2100
ttcaagcat	gagaagccgc	ccaggagaaa	atttagtctt	tcattgagcc	caaatcagct	2160
ctctctgtgc	ctgagctctt	tgacacatgt	gaactctcta	tttaggacata	caaaagagga	2220
gagagtgagg	gcggagagag	agagaggaga	gagaacacga	acgattctcta	gtctgaggtt	2280
ctgtgagccg	cctactcttt	ttgaaaatct	cacatgggtc	aggaagcagg	gaggagaaac	2340
caagtcttag	gtttacactt	tgaatctcca	agttgcata	tataaaggga	tattgtatca	2400
gccaaagtgc	attaataaca	ggaagcagga	tattctattg	caatgggtgc	ctatttggtta	2460
tctctgtctc	caggctctct	cccaccagtc	tgctctctac	caggatcaga	gtgatgttcc	2520
tacagtgtcc	ctgtctctgt	ctctctctgt	cctgaagctc	ttaattggcc	ccatgtctacc	2580
cacaagccag	aggtccagct	ccttgacatg	atcagaagga	tcttcatcat	gagaccaccag	2640
gtgctctctc	aacccctgtg	gcattccctt	cctgttgtaa	tgtatatatt	acattccagc	2700
aatattgcac	ctattttaca	tttgccaagc	actccatgct	atctttatgt	ccatcccttg	2760
gcatactgt	atctttctac	tggatttttt	tttttcccat	tactccattg	catatgcttc	2820
tttgaactctg	gcataattgt	ctctctgtgt	catcaccttg	caattttatt	gtcatgtctt	2880
gtcttctcta	ccagactatg	agctgcttgg	tgcaaggact	gcgagttatt	catcactgtg	2940
gccctatgcc	tggtagagca	tcagtacctta	gaaggcactc	agcctgattt	tgtgggggtga	3000
atggatgggt	ggatggatga	cgagagtctt	acaagagaaa	ttgggataggt	ttgggacaa	3060
atggttaagt	tatccatgta	acagaccccc	agagaagaca	acaaatggcc	tcttctgtaa	3120
agctcagaat	ctgagagatg	ggagtaagcc	agacaaggta	tctagtccag	aataggggaag	3180
ttgggatgat	atggtgacat	gctgtgggac	tgacttctct	ttctctctag	ataagagccc	3240
ttggagagac	aggcagccag	aagcacctgt	gctcccagga	taagggtgag	catcaggatg	3300
gactgtggag	aaggagccgc	ctgatgcgca	cttccactgtg	gacaaaacga	acatctccct	3360
ctggccccga	gcagccacaca	tcgtgtgtga	gaacctgtct	cggtgtctgt	gtgcaaacct	3420
gcccttttgt	gctctctcaa	cacacatttt	cttctctctc	caacagagctt	tctcccacag	3480
tcgggtccat	ctctgtcttc	ggggaaaaaca	cccacagctc	gtgctgcatt	aatctgctgt	3540
acgctgtctc	tggtcgtctc	cgctcgtctc	caggcgtctc	tttgaagtc	ctcatgttct	3600
atgctctggg	cttagccctc	ctctgtgcca	gccggctctc	ttcagatgca	gaccactctc	3660
ctgctctccg	ggttctctct	cctgtggctt	tttcatttgt	ctctctctc	cttcttccat	3720
gtgcatgaac	aggtctgtgt	gccccagta	cggtgagctg	atgctcttct	cttcccattt	3780
ctctggagat	tattgggatt	agcatttgca	cattggtgct	caaggatata	gtctttgtct	3840
tcaggaaaca	ttgatctagt	ttacatctct	gtgctatttt	ctcccgtctc	acccccacc	3900
aatctgagct	ctgctcctgt	gatctagagc	cttgtggaca	gtcttgata	gtgcatgacc	3960
tctgaccct	gtgaaggacc	gagcctcatg	tatcatctgc	ccagctcac	gcagtgaga	4020
gcctggccaa	cgagccaaaca	gatctccatg	actcagctcc	gtctccccag	gactcgttgg	4080
tgctgtcttc	attctctctg	ccctgtgaag	atcacatcta	ccgagctgtc	gtctcatttc	4140
gatgttgcat	ggttttatct	tcttctcttt	cttgctctgt	caggcttcat	ggccttttgt	4200
ctggccagag	ctctctctct	cctggccagg	ctccagcaag	cagagcaaa	ggaccacaa	4260
agctgttgag	tttttttttc	tctctctgtg	gcctcgggac	atatttgggc	ttagacttgt	4320
agggcttgag	cagagctccc	cttgccccat	cctagacccc	tgccctctaa	catcgctatt	4380
tctccaggcc	tgccctttga	gcctcttggt	ttaactttga	tctctctctc	ggaccacag	4440
cttgggacct	gagcagaatg	tgaaagtggg	ttggggcaag	gaagggggaga	aacagttttg	4500
taatctcttc	ttccatgttc	ttctggagaag	ccacttccag	attagtggct	ggttctctcc	4560
atggtccacg	aggggccccat	ggacagatgt	ggggggagtg	tgctgtctcta	gcagatggcc	4620
actgcagggg	tttctgaaaa	cagagggatg	gcaaccaagg	gtgggggtct	agggggaaat	4680
aagttctggg	gacagtgggt	gggctcatgg	agggcaccct	ttatcaaatg	ttctctgaac	4740
actcagaaaa	ttcaggaatg	gtttcttaact	ctgctctgtg	ctgtcctgtg	aaatcttttc	4800
acagagagcc	tgtgttttat	caatctcccc	attatctgta	tcacactgtg	ttcttggaac	4860
atgggttagtg	cccatgtcac	gtttgttgtta	cgttaatgta	tgtattggag	gttgggggtg	4920
cccatctggac	tgtcttgggt	ctttgggaag	cttcagcctca	ttcttctctc	tctcttgatc	4980
aaactgcacaa	cacccecatc	cctgtccctg	ggactccctc	cagctcagct	ctctgatttc	5040
tcaatccagc	atccccgggt	tatgggcacc	atatcagatg	taaaagacaa	gttccagttg	5100
ctgaaaggct	gtgtggacaa	catcagcacc	ctggattctg	aaatttttaa	gaatagtgac	5160
ggcatggagg	cagctggcgt	tcagatccag	atggtgaatg	agagcctggg	ttatgtgcgt	5220

tctcagttcc	tgaagttaaa	aaccagtggt	gagaaggcca	acgcacagat	ccagatctta	5280
acaagaagtt	gggaagaagt	cagttaccctta	aatgcgccaaa	tcccagagtt	aaaaagtgat	5340
ttggagaaag	ccagtgcttt	aaatacaaaag	atccggggcac	tccaggggcag	cttggagaat	5400
atgagcaagt	tgctcaaacg	acaaagttaag	tgactcagaa	aattacattg	aagctgcacca	5460
gtggcccatg	ggatctctacc	tgtcccagac	ctgaggccat	tggtctgggt	ggttggggag	5520
gagagtgggg	gcaaaagagag	ggcagccatg	ggctaggaaag	ttaaggagag	agggcttggag	5580
gttggggag	acttaggggc	tgttaggaga	aaagagacca	gggtccagag	agagctccga	5640
ccacaaaagt	cagaattgtaa	aagcatttag	ggatgtccac	cctggcccac	acctagtcac	5700
ttccctcaaa	gtctctctct	agagtcacag	ggctcagcca	cttgtcatgg	ccgattggag	5760
gttgcttctc	catcatgggg	aagacactct	ttgtcccaacc	tcttgcatta	taacctctcc	5820
agtcccagag	actctattag	tctctgtctg	actttcagga	tttgaaggag	gttccctaat	5880
ctcctataca	aggacccaag	gacaccacg	cacagctcca	tttgctgctg	tctctgagag	5940
ctcattcaag	tgccccccac	aagccagcat	cccaagaata	caagataacc	agcgttcact	6000
tttacctctt	gtctctctaga	tgatattcta	caggtgggtt	ctcaaggctg	gaagtacttc	6060
aaggggaaat	tctattactt	ttctctcatt	ccaaagacct	ggtatagtgc	cgagcagttc	6120
tggtgtgtcca	ggaattccaca	cctgacctcg	gtgacctcag	agagtgcaga	ggtgagtgtc	6180
gtgctctatg	gctctgtgaa	gggggcgtat	gagcacttgg	ccaggagaga	tgggcaagat	6240
tatacttcgt	gaacaaaata	ccccaaata	tgatgcagta	atgaagaaag	attgtgtctc	6300
agtagcatgt	cgaataaggg	cgggcaagg	ggctgtgctc	actgtggcta	ctcaaggacc	6360
caggcccaag	gagttctcat	cttaatatgt	ctccacaatt	ctgtggggcag	gaaaggggaa	6420
cttgacatat	tgtgcgaagc	ttctgccccag	atttaacata	cctcattttt	gcttacattt	6480
cactagctta	agttatgtta	attaagttaa	gtaagttatg	cagtaactct	ggattgggag	6540
gatcctggat	tcagtagtca	ctggaacagg	gcagagagga	tctactttcc	atgtgcccag	6600
aaaaaaagaa	atatttgtga	acagccctaa	tgattccaca	tgactcaaga	agcttccctg	6660
ctggtgaggg	agaaatggg	caggcccttt	tcactctggga	ggtgggagat	cagagcaggt	6720
cagagccctg	gctctggcgt	agttctagag	cctgaacctt	gccattcaac	tagtccctag	6780
agctctgggt	gaatacccca	cttcaactgg	caaaactcac	tggtccctca	ttgatgaaac	6840
atgcatgtgt	atggcacctg	cctcagagga	aaggagagaa	tgcatatgga	cagatcagca	6900
cagtgccgtg	ttgtggattca	gggtctcatt	aattcaggta	ttatcatatg	aaccttcttc	6960
ttgctggctg	tgctctggag	ttcagctgag	gccttctgtg	gcttcagcac	ctgcttctct	7020
agtgccagaa	aggcttgagt	cctgagcttg	ttagctgcag	agcagggaca	catcataatc	7080
tggaagatga	aaactctggct	gtgggcaagg	gcaggaaagaa	gcttgagagg	ccagtttgtg	7140
cagcgcatct	gtgggtcagg	gcgtgtcact	agcgccaggt	aagaacaccc	agagacagat	7200
gatcaagctc	caagtgtggc	cgcacctctg	cttatctctg	ctttctcaca	ggagttctct	7260
tataaaacag	cggggggaact	catctacttg	attggcctga	ctaaagcagg	gttggaagg	7320
ggatgtctct	gggtggatga	cacgccattc	aacaaggctc	aaagtgtgag	gatagccctc	7380
ggagccctcc	gtgcagacct	gactttcccc	ggccatggcc	agggatgag	gggagtggg	7440
gcgatgttcc	ccatgagaca	gggtttctga	ttcttccctg	tcttagagt	acaggaatgc	7500
tgcaaccaag	atccagacaca	accctgtcac	caactggctg	tggaactgag	ccctccacgc	7560
ctctgtgggt	ttggcaacaa	ggcctctcac	ctggccagct	tcagggatct	gtcttagagt	7620
ctaggtcttc	acagtggtgg	tttgtgtagg	gaactgaaag	tggtgggtgt	gttgggctg	7680
gacttggggc	atgtgaaagg	ttagaggtcg	aaagtccaac	agtcctcttc	ccttaggctt	7740
tggtgtcaga	catttgcctat	gccattccct	aggacaaaag	cttgggttga	gttaactcat	7800
ttcttctact	gaaataagtt	ctttttgatt	ttccactttg	taaaatccat	gttttcccg	7860
ctcttggtag	gttctgtgatt	ccaggtgagc	cttggggagc	ccaacaatgc	gaacactgtg	7920
gcaataaaaa	ggctccctca	cttcaggcct	ggaatgctg	cccatgtgac	aaaaagcttc	7980
ttttcatttg	taagcgaccc	tatgtcccat	cagaaacagc	acagtgacag	ctccaaagct	8040
cactctttga	gtcccaacgc	ttgttaaaaa	tgaggaaatg	cctctttctt	ccccagactc	8100
caggatgact	ttgcacgtta	atttttctgt	cttcaaaatt	gtcccaagct	ggcatctcgt	8160
agtcogtctg	tcttgggtctg	aaattctctg	agctcttggg	ggcagcttga	atgggaactga	8220
gaattcagg	taaaagtggg	ggggtgggta	gagaggattt	agaagtttca	atgtccctgc	8280
taaggaggat	caagaccctg	aatccggcct	aacaccctgg	gggttttccac	tcttccagag	8340
aaacctcagc	ttcatcacat	caaagttact	ccagagcaac	caagcaattc	ctctgatatt	8400
gcctatccag	gctttttctg	gccaaaaccc	ctagaatttc	catgtctctg	cttagctgtg	8460
ctggcagcta	gcagcttgct	gtgttttcag	tgcaaatatg	tctgttcttg	gaaactctgc	8520
tcatggtatg	tccccagtg	tttcttcac	cacatcatct	aaagcctgaa	ctgtctcttc	8580
cttggtttcaa	gtcagttgct	gacacggact	tgtatctctc	ctcagagctc	gcttgcaacc	8640
agctccctct	ctcccttcac	ttccttagta	cactggagtg	ccagccctg	cttccaacct	8700
agctccctct	cagccctctg	cctcacctct	ccggcacctc	ctctctcttc	tgcatattct	8760
atcttctcgt	gtcttctgta	tggggaagcag	cttccagtgc	tcacttgaa	tcaccttcca	8820
gcttctctcag	aataaaatgc	tgctctgggtc	aaggactcac	tccaagtga	cttttctcatt	8880

09787192.031501

```

tctggttgtc cagggtgaata tgtgggaaag gcagctctcct ctggttgaca tgaagttcta 8940
gggtatctctc aggaataatc tggggagtc aaaaatacaa ggactgggga agttccagtc 9000
ctggaaatgc cacaataatgt gaccagctact tatctctagt ttttattaaa gtagagcaag 9060
gtctccaatg tcacgatctt ggtgatcttt ctctctgttt actgcacaa ctcttagtct 9120
atagctcaat tcccaagaac aagtctcagc aggttcccca ctcttcacag agaccagttt 9180
ccacaggcat cagttccaaa tcccaagtc agtggtctga gctggaatcc aggcagcagc 9240
caccacagag agggagaggag ggtggagtga gcacaggctc tcattaaagg cctcaggaaa 9300
agatgctccc ttaataaact gtaaccagca gtgtgttgtt ctgggtgc aaatgggtcaca 9360
gctgaggcca caggcttgta accataccctc gagtctccctg cctctcacc aatataatct 9420
tgcttcacag cggcccaaaag acctaaacctc atcatctgga ccagctgcgat gccaccagtc 9540
tgccctcggt ccccacctgg ctaatttctc agcctgagaa actattcaaa caagccaatc 9600
ttctacttta atgggtttca ctctctctgc atccttttat tactacaagg cctgctcccc 9660
acatctctct acaggaatcc ggggcatctc atccttttat tactacaagg cctgctcccc 9720
acagcctctg ctggttcaact ctgctcctga ggggtgaccc atgtggccctc gtgtggctta 9780
tgatatcttt cccacagtaga ctgtatttgt gactagtaaa ctgctgccag tctcactctg 9840
acagtgtaaa atgtcttgtt ttggccatct tgtcctattt agagcagggg atccctcccc 9900
caccatggga ggaatggga ggtgacaaga acaaggctgg tcggggatct ctggttggtt 9960
ttggcaaaag gatgagctgg gaaatcaga ccatttctct gggaaagaat ttgaaccagg 9960
aaatagcaag aggatgaggc tgtaacaaa aggaagtga gctggaaggc actgagttaa 10020
gagaaaggct ggagggggcc tcacgtggca ttggaagaaa ctagcaatga gcagaagcta 10080
tgaggcaggg gaaagacatc aatacaggcc tggcatggtg gctcatgctc ataataccag 10140
caccttggga ggtcgaagtg ggtgggtcac ctgaggtcag gagttcgaga cagcctggcc 10200
aacatggtag aaccccatct ctactaaaaa agccggggat agccggggat ggtggtgggc 10260
acctgttaatt ccaactactt gggagactga ggcaggagaa ttgcttgaat ctgggaggca 10320
gaggttgtag tgagctgaga tcccaccact gcactccagc ctggggcaaca gggcaagact 10380
ttgtttcaaa aaaaagaag tgactgcaga ggattatagt tggcagagaa aagagaacgg 10440
ctcagaggag tcgcaatgga ggtcccggag ggcagcctga agggctccgg ctgctcccg 10500
tcccagggtc gctcagatc ctcccagccc ttctgatctc cctggtttct gtgcatgggg 10560
acctacgag gctgtgctcc tgaccccaac cattgtcttt tctgaaact gaaagagcct 10620
gagtcagtga ggaatgtgtt ttatctggag tctgtgcccc agc 10663

```

```

<210> 4
<211> 120
<212> DNA
<213> homo sapiens

```

```

<400> 4
ggcagccaga agcactctgt ctcccaggat aagggtgagc actcaggatg actgtggaga 60
aggaggcccc tgatgcgcac ttactctgtg acaaacagaa catctccctc tggcccccag 120

```

```

<210> 5
<211> 117
<212> DNA
<213> homo sapiens

```

```

<400> 5
agctctctcc caagtcctgt ccatctctgt tcccggggaa aacacccaca gtccgtgtgtg 60
cattaatctg cctgacgtgt gtctctgtgt cctccgtctc gctgcaggcc gtccctt 117

```

```

<210> 6
<211> 375
<212> DNA
<213> homo sapiens

```

```

<400> 6

```

```

atccccgggt tatgggcacc atatcagatg taaagaccaa tgtccagttg ctgaaaggtc 60
gtgtggacaa catcagcacc ctggattctg aaattaaaaa gaatagtac ggcatggagg 120
cagctggcgt tcagatccag atggtgaatg agagcctggg ttatgtgcgt tctcagttcc 180
tgaagttaaa aaccagtggt gagaaggcca acgcacagat ccagatctta acaagaagtt 240
gggaagaaat cagtacctta aatgccccaa tccagagttt aaaaagtgat ttggagaaaag 300
ccagtgcctt aaatacaaa atccgggcac tccagggcag cttggagaat atgagcaagt 360
tgctcaaacg acaaa
375

```

```

<210> 7
<211> 152
<212> DNA
<213> homo sapiens

```

```

<400> 7
atgatattct acaggtgtgtt tctcaaggct ggaagtactt caaggggaac ttctattact 60
tttctctcat tccaaagacc tggatatgtg ccgagcagtt ctgtgtgtcc aggaattcac 120
acctgacctc ggtgacctca gagagtggag ag
152

```

```

<210> 8
<211> 119
<212> DNA
<213> homo sapiens

```

```

<400> 8
gagtttctgt ataaaacagc ggggggactc atctactgga ttggcctgac taaagcaggg 60
atggaagggg actggtctctg ggtggatgac acgccattca acaaggtcca aagtgcgag 119

```

```

<210> 9
<211> 985
<212> DNA
<213> homo sapiens

```

```

<400> 9
gttctggatt ccaggtggagc ccaacaatgc tgggaacaat gaacactgtg gcaatataaa 60
ggctccctca ctccaggcct ggaatgatgc cccatgtgac aaaacggttc ttctcatttg 120
taagcgacc tcagtcccat cagaaccctg acaggacagg ctcccaagct cactctttga 180
gtcccaacgc ttgttaaaac tgaggaaatg cctctttctt ccccgagact caggatgact 240
ttgcacgtta atttttcttg ctccaaaaat gtcccacagt ggcattctcg agtccgctcg 300
tcttggctgg aaattctctg acgtctctga ggcagctgga atggaaaagg gaattcagg 360
taagttggga ggggtgggta gagaggattt agaagttcca attgccctgc taaggaggat 420
caagaccctg aatccggcac aacaccctgg ggttttccac tcttccagag aaacctcagc 480
ttcatcacat caaagttaact ccagagcaac caagcaattc tcttgatatt gtcatccagg 540
gcttttcttg gccaaacccc ctagaatttc catgtctctg cttagctgtg ctggcagcta 600
gcagctggct gtgttttcag tgcataatgc tctgttcttg gaaatccctg tcatggtgatg 660
tcccagtggt ttcttctcag cacatcatct aaagcctgaa cccgttcttc tctggttcaa 720
gtcagtggtt gacacggact tgtatctctc tcagagctcg gctggcacc accctccctt 780
ctctctccac tcccttagta cactggagtg ccgagccctg ccttccacc agcgtccatc 840
cagccctctg cctcacctct ccggcacctc ctctcctctc tgcatttctt atcttctctg 900
gtcttggtca tgggaagctg ccttcagtcg ctctcatgat tcaccttcca gcttccctag 960
aataaaatgc tgcctgggtc aagga
985

```

400> 10	ggcctctttg	ctcgtttggg	gtcttcagtt	aagatgatta	aaatgtctgt	gtaccagggg	60										
	agtagtcagg	tggcttcttc	atcaaaagccc	aattcttagt	caaaaggatg	aagctgtggc	120										
	ctctgcctgg	ctaatcgtct	aagccaagaa	ctgggaacct	gggcatgaca	aagtggcctg	180										
	ctttttggct	catttcccg	tctttcttgg	atgaaaaggt	ccttggggag	acggatatct	240										
	agcttttctc	aagccagagg	cagag	atg	ttg	gag	gag	gct	ccc	gaa	gcg	cac	292				
				Met	Leu	Glu	Glu	Ala	Pro	Glu	Ala	His					
				1					5								
	ttc	aca	gtg	gac	aaa	cag	aac	atc	tct	ctc	tgg	cct	cga	gag	cct	cct	340
	Phe	Thr	Val	Asp	Lys	Gln	Asn	Ile	Ser	Leu	Trp	Pro	Arg	Glu	Pro	Pro	
	10					15					20					25	
	ccc	aag	caa	gat	ctg	tct	cca	ggt	ctg	agg	aaa	cct	ctc	tgt	atc	tgc	388
	Pro	Lys	Gln	Asp	Leu	Ser	Pro	Val	Leu	Arg	Lys	Pro	Leu	Cys	Ile	Cys	
					30					35					40		
	gtg	gcc	ttc	acc	tgc	ctg	gca	ttg	gtg	ctg	gtc	acc	tcc	att	gtg	ctt	436
	Val	Ala	Phe	Thr	Cys	Leu	Ala	Leu	Val	Leu	Val	Thr	Ser	Ile	Val	Leu	
					45				50					55			
	cag	gct	ggt	ttc	tat	cct	agg	ttg	atg	ggc	aaa	ata	ttg	gat	gtg	aag	484
	Gln	Ala	Val	Phe	Tyr	Pro	Arg	Leu	Met	Gly	Lys	Ile	Leu	Asp	Val	Lys	
					60			65					70				
	agt	gat	gcc	cag	atg	ttg	aaa	ggt	cgt	gtg	gac	aac	atc	agc	acc	ctg	532
	Ser	Asp	Ala	Gln	Met	Leu	Lys	Gly	Arg	Val	Asp	Asn	Ile	Ser	Thr	Leu	
		75					80					85					
	ggt	tct	gat	ctt	aag	act	gaa	aga	ggt	cgt	gtg	gac	gat	gct	gag	ggt	580
	Gly	Ser	Asp	Leu	Lys	Thr	Glu	Arg	Gly	Arg	Val	Asp	Asp	Ala	Glu	Val	
	90					95					100					105	
	cag	atg	cag	ata	gtg	aac	acc	acc	ctc	aag	agg	gtg	cgt	tct	cag	atc	628
	Gln	Met	Gln	Ile	Val	Asn	Thr	Thr	Leu	Lys	Arg	Val	Arg	Ser	Gln	Ile	
					110					115					120		
	ctg	tct	ttg	gaa	acc	agc	atg	aag	ata	gcc	aat	gat	cag	ctc	ctg	ata	676
	Leu	Ser	Leu	Glu	Thr	Ser	Met	Lys	Ile	Ala	Asn	Asp	Gln	Leu	Leu	Ile	
				125					130					135			
	tta	aca	atg	agc	tgg	gga	gag	ggt	gac	agt	ctc	agt	gcc	aaa	atc	cca	724
	Leu	Thr	Met	Ser	Trp	Gly	Glu	Val	Asp	Ser	Leu	Ser	Ala	Lys	Ile	Pro	
			140					145					150				

gaa ctg aaa aga gat ctg gat aaa gcc agc gcc ttg aac aca aag gtc 772
 Glu Leu Lys Arg Asp Leu Asp Lys Ala Ser Ala Leu Asn Thr Lys Val
 155 160 165

caa gga cta cag aac agc ttg gag aat gtc aac aag ctg ctc aaa caa 820
 Gln Gly Leu Gln Asn Ser Leu Glu Asn Val Asn Lys Leu Leu Lys Gln
 170 175 180 185

cag agt gac att ctg gag atg gtg gct cga ggc tgg aag tat ttc tcg 868
 Gln Ser Asp Ile Leu Glu Met Val Ala Arg Gly Trp Lys Tyr Phe Ser
 190 195 200

ggg aac ttc tat tac ttt tca cgc acc cca aag acc tgg tac agc gca 916
 Gly Asn Phe Tyr Tyr Phe Ser Arg Thr Pro Lys Thr Trp Tyr Ser Ala
 205 210 215

gag cag ttc tgt att tct aga aaa gct cac ctg acc tca gtg tcc tca 964
 Glu Gln Phe Cys Ile Ser Arg Lys Ala His Leu Thr Ser Val Ser Ser
 220 225 230

gaa tgg gaa caa aag ttt ctc tac aag gca gca gat gga att cca cac 1012
 Glu Ser Glu Gln Lys Phe Leu Tyr Lys Ala Ala Asn Gly Ile Pro His
 235 240 245

tgg att gga ctt acc aaa gca ggg agc gaa ggg gac tgg tac tgg gtg 1060
 Trp Ile Gly Leu Thr Lys Ala Gly Ser Glu Gly Asp Trp Tyr Trp Val
 250 255 260 265

gac cag aca tca ttc aac aag gag caa agt agg agg ttc tgg att cca 1108
 Asp Gln Thr Ser Phe Asn Lys Glu Gln Ser Arg Arg Phe Trp Ile Pro
 270 275 280

ggt gaa ccc aac aac gca ggg aac aac gag cac tgt gcc aat atc agg 1156
 Gly Glu Pro Asn Asn Ala Gly Asn Asn Glu His Cys Ala Asn Ile Arg
 285 290 295

gtg tct gcc ctg aag tgc tgg aac gat ggt ccc tgt gac aat aca ttt 1204
 Val Ser Ala Leu Lys Cys Trp Asn Asp Gly Pro Cys Asp Asn Thr Phe
 300 305 310

ctt ttc atc tgc aag agg ccc tac gtc caa aca act gaa tgacagatct 1253
 Leu Phe Ile Cys Lys Arg Pro Tyr Val Gln Thr Glu
 315 320 325

ggccctgagct cggcatctgt ggggcaacag tgacctggct gaagagatgt ctctctccct 1313

gaggetccaa gattgctctg tacttacgtt tttttcttgc ttgaaaattg tcccaaacac 1373

agcctgtgggt ctttctgtct tggctggcag ttctctgtct ctggaggcct tggaggagct 1433

tgggttaaac ggggtaggac ctgaaaaggg tgtagcagtc cttactgccc aggcgaggca 1493

ggtcagcaca ccaaacaggt tgtttagatt ttctctgatcc ttctcagaag ccttggctga 1553

ccataataaa gctacattca aatatgacca gtatttgagg agacagacat gcccaaat 1613

aaccatgata caatttatcc aacatgtatt agaacacctc atggtagtt caaaatagta 1673

aatatgttgt ttttatgtgc ctattgcaaa taaatgtaaa gacttaaaaa aaaaaaaaaa 1733

aaaaaaaaaa aaaaaaaaaa aaa 1756

09787192-031501

<210> 11
 <211> 326
 <212> PRT
 <213> Mus musculus

<400> 11
 Met Leu Glu Glu Ala Pro Glu Ala His Phe Thr Val Asp Lys Gln Asn
 1 5 10 15
 Ile Ser Leu Trp Pro Arg Glu Pro Pro Lys Gln Asp Leu Ser Pro
 20 25 30
 Val Leu Arg Lys Pro Leu Cys Ile Cys Val Ala Phe Thr Cys Leu Ala
 35 40 45
 Leu Val Leu Val Thr Ser Ile Val Leu Gln Ala Val Phe Tyr Pro Arg
 50 55 60
 Leu Met Gly Lys Ile Leu Asp Val Lys Ser Asp Ala Gln Met Leu Lys
 65 70 75 80
 Gly Arg Val Asp Asn Ile Ser Thr Leu Gly Ser Asp Leu Lys Thr Glu
 85 90 95
 Arg Gly Arg Val Asp Asp Ala Glu Val Gln Met Gln Ile Val Asn Thr
 100 105 110
 Thr Leu Lys Arg Val Arg Ser Gln Ile Leu Ser Leu Glu Thr Ser Met
 115 120 125
 Lys Ile Ala Asn Asp Gln Leu Leu Ile Leu Thr Met Ser Trp Gly Glu
 130 135 140
 Val Asp Ser Leu Ser Ala Lys Ile Pro Glu Leu Lys Arg Asp Leu Asp
 145 150 155 160
 Lys Ala Ser Ala Leu Asn Thr Lys Val Gln Gly Leu Gln Asn Ser Leu
 165 170 175
 Glu Asn Val Asn Lys Leu Leu Lys Gln Gln Ser Asp Ile Leu Glu Met
 180 185 190
 Val Ala Arg Gly Trp Lys Tyr Phe Ser Gly Asn Phe Tyr Tyr Phe Ser
 195 200 205
 Arg Thr Pro Lys Thr Trp Tyr Ser Ala Glu Gln Phe Cys Ile Ser Arg
 210 215 220
 Lys Ala His Leu Thr Ser Val Ser Ser Glu Ser Glu Gln Lys Phe Leu
 225 230 235 240
 Tyr Lys Ala Ala Asp Gly Ile Pro His Trp Ile Gly Leu Thr Lys Ala
 245 250 255
 Gly Ser Glu Gly Asp Trp Tyr Trp Val Asp Gln Thr Ser Phe Asn Lys
 260 265 270

00787102-031501
 105150-2612860

Glu	Gln	Ser	Arg	Arg	Phe	Trp	Ile	Pro	Gly	Glu	Pro	Asn	Asn	Ala	Gly
		275					280					285			
Asn	Asn	Glu	His	Cys	Ala	Asn	Ile	Arg	Val	Ser	Ala	Leu	Lys	Cys	Trp
		290					295					300			
Asn	Asp	Gly	Pro	Cys	Asp	Asn	Thr	Phe	Leu	Phe	Ile	Cys	Lys	Arg	Pro
		305				310					315				320
Tyr	Val	Gln	Thr	Thr	Glu										
				325											

09787192.031501

SEQUENCE LISTING

<110> Schering Corporation

<120> Antibodies to Mammalian Langerhans Cell Antigen and Their Uses

<130> SF0820K2 PCT

<140> PCT/US99/22269

<141> 1999-09-23

<150> EP 99 400 394.5

<151> 1999-02-18

<150> EP 98 402 374.7

<151> 1998-09-25

<160> 11

<210> 1

<211> 1547

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (56) .. (1039)

<400> 1

ctggaaaggg cagccagaag cacctgtgct cccaggataa ggggtgagcac tcagg atg 58
 Met
 1

act gtg gag aag gag gcc cct gat gcg cac ttc act gtg gac aaa cag 106
 Thr Val Glu Lys Glu Ala Pro Asp Ala His Phe Thr Val Asp Lys Gln
 5 10 15

aac atc tcc ctg tgg ccc cga gag cct cct ccc aag tcc ggt cca tct 154
 Asn Ile Ser Leu Trp Pro Arg Glu Pro Pro Pro Lys Ser Gly Pro Ser
 20 25 30

ctg gtc ccg ggg aaa aca ccc aca gtc cgt gct gca tta atc tgc ctg 202
 Leu Val Pro Gly Lys Thr Pro Thr Val Arg Ala Ala Leu Ile Cys Leu
 35 40 45

acg ctg gtc ctg gtc gcc tcc gtc ctg ctg cag gcc gtc ctt tat ccc 250
 Thr Leu Val Leu Val Ala Ser Val Leu Leu Gln Ala Val Leu Tyr Pro
 50 55 60 65

cgg ttt atg ggc acc ata tca gat gta aag acc aat gtc cag ttg ctg 298
 Arg Phe Met Gly Thr Ile Ser Asp Val Lys Thr Asn Val Gln Leu Leu
 70 75 80

aaa ggt cgt gtg gac aac atc agc acc ctg gat tct gaa att aaa aag 346
 Lys Gly Arg Val Asp Asn Ile Ser Thr Leu Asp Ser Glu Ile Lys Lys
 85 90 95

aat agt gac ggc atg gag gca gct ggc gtt cag atc cag atg gtg aat 394
 Asn Ser Asp Gly Met Glu Ala Ala Gly Val Gln Ile Gln Met Val Asn
 100 105 110

09787192-0345004

gag agc ctg ggt tat gtg cgt tct cag ttc ctg aag tta aaa acc agt Glu Ser Leu Gly Tyr Val Arg Ser Gln Phe Leu Lys Leu Lys Thr Ser 115 120 125	442
gtg gag aag gcc aac gca cag atc cag atc tta aca aga agt tgg gaa Val Glu Lys Ala Asn Ala Gln Ile Gln Ile Leu Thr Arg Ser Trp Glu 130 135 140 145	490
gaa gtc agt acc tta aat gcc caa atc cca gag tta aaa agt gat ttg Glu Val Ser Thr Leu Asn Ala Gln Ile Pro Glu Leu Lys Ser Asp Leu 150 155 160	538
gag aaa gcc agt gct tta aat aca aag atc cgg gca ctc cag gcc agc Glu Lys Ala Ser Ala Leu Asn Thr Lys Ile Arg Ala Leu Gln Gly Ser 165 170 175	586
ttg gag aat atg agc aag ttg ctc aaa cga caa aat gat att cta cag Leu Glu Asn Met Ser Lys Leu Leu Lys Arg Gln Asn Asp Ile Leu Gln 180 185 190	634
gtg gtt tct caa ggc tgg aag tac ttc aag ggg aac ttc tat tac ttt Val Val Ser Gln Gly Trp Lys Tyr Phe Lys Gly Asn Phe Tyr Tyr Phe 195 200 205	682
tct ctc att cca aag acc tgg tat agt gcc gag cag ttc tgt gtg tcc Ser Leu Ile Pro Lys Thr Trp Tyr Ser Ala Glu Gln Phe Cys Val Ser 210 215 220 225	730
agg aat tca cac ctg acc tcg gtg acc tca gag agt gag cag gag ttt Arg Asn Ser His Leu Thr Ser Val Thr Ser Glu Ser Glu Gln Glu Phe 230 235 240	778
ctg tat aaa aca gcg ggg gga ctc atc tac tgg att ggc ctg act aaa Leu Tyr Lys Thr Ala Gly Gly Leu Ile Tyr Trp Ile Gly Leu Thr Lys 245 250 255	826
gca ggg atg gaa ggg gac tgg tcc tgg gtg gat gac acg cca ttc aac Ala Gly Met Glu Gly Asp Trp Ser Trp Val Asp Asp Thr Pro Phe Asn 260 265 270	874
aag gtc caa agt gcg agg ttc tgg att cca ggt gag ccc aac aat gct Lys Val Gln Ser Ala Arg Phe Trp Ile Pro Gly Glu Pro Asn Asn Ala 275 280 285	922
ggg aac aat gaa cac tgt ggc aat ata aag gct ccc tca ctt cag gcc Gly Asn Asn Glu His Cys Gly Asn Ile Lys Ala Pro Ser Leu Gln Ala 290 295 300 305	970
tgg aat gat gcc cca tgt gac aaa acg ttt ctt ttc att tgt aag cga Trp Asn Asp Ala Pro Cys Asp Lys Thr Phe Leu Phe Ile Cys Lys Arg 310 315 320	1018
ccc tat gtc cca tca gaa ccg tga c aggacaggct cccaagctca Pro Tyr Val Pro Ser Glu Pro 325	1063
ctctttgagc tccaacgctt gttaaacatg aggaatgcc tctttcttcc ccagactcca	1123
ggatgacttt gcacgttaat ttttcttgct tcaaaattgt cccacagtgg cattctggag	1183
tccgtctgtc ttggctggaa attctctgac gtcttgagg cagctggaat ggaaaggaga	1243

attcaggtta aagtgagg ggtgggtaga gaggatttag aagtccaat tgcctgcta 1303
 aggagagatca agaccgtaa tccggcacia caccctgggg ttttccactc ttcagagaa 1363
 acctcagctt catcacatca aagttactcc agagcaacca agcaattctc ctgatattgt 1423
 catccagggc ttttcttgcc caaaccacct agaatttcca tgtctctgct tagctgtgct 1483
 ggcagctagc agctggctgt gtttcagtg caaatagctc tgttcttgga aatcctgctc 1543
 atgg 1547

<210> 2

<211> 328

<212> PRT

<213> Homo sapiens

<400> 2

Met Thr Val Glu Lys Glu Ala Pro Asp Ala His Phe Thr Val Asp Lys
 1 5 10 15
 Gln Asn Ile Ser Leu Trp Pro Arg Glu Pro Pro Pro Lys Ser Gly Pro
 20 25 30
 Ser Leu Val Pro Gly Lys Thr Pro Thr Val Arg Ala Ala Leu Ile Cys
 35 40 45
 Leu Thr Leu Val Leu Val Ala Ser Val Leu Leu Gln Ala Val Leu Tyr
 50 55 60
 Pro Arg Phe Met Gly Thr Ile Ser Asp Val Lys Thr Asn Val Gln Leu
 65 70 75 80
 Leu Lys Gly Arg Val Asp Asn Ile Ser Thr Leu Asp Ser Glu Ile Lys
 85 90 95
 Lys Asn Ser Asp Gly Met Glu Ala Ala Gly Val Gln Ile Gln Met Val
 100 105 110
 Asn Glu Ser Leu Gly Tyr Val Arg Ser Gln Phe Leu Lys Leu Lys Thr
 115 120 125
 Ser Val Glu Lys Ala Asn Ala Gln Ile Gln Ile Leu Thr Arg Ser Trp
 130 135 140
 Glu Glu Val Ser Thr Leu Asn Ala Gln Ile Pro Glu Leu Lys Ser Asp
 145 150 155 160
 Leu Glu Lys Ala Ser Ala Leu Asn Thr Lys Ile Arg Ala Leu Gln Gly
 165 170 175
 Ser Leu Glu Asn Met Ser Lys Leu Leu Lys Arg Gln Asn Asp Ile Leu
 180 185 190
 Gln Val Val Ser Gln Gly Trp Lys Tyr Phe Lys Gly Asn Phe Tyr Tyr
 195 200 205
 Phe Ser Leu Ile Pro Lys Thr Trp Tyr Ser Ala Glu Gln Phe Cys Val
 210 215 220

09787192-031501

Ser Arg Asn Ser His Leu Thr Ser Val Thr Ser Glu Ser Glu Gln Glu
225 230 235 240

Phe Leu Tyr Lys Thr Ala Gly Gly Leu Ile Tyr Trp Ile Gly Leu Thr
245 250 255

Lys Ala Gly Met Glu Gly Asp Trp Ser Trp Val Asp Asp Thr Pro Phe
260 265 270

Asn Lys Val Gln Ser Ala Arg Phe Thr Ile Pro Gly Glu Pro Asn Asn
275 280 285

Ala Gly Asn Asn Glu His Cys Gly Asn Ile Lys Ala Pro Ser Leu Gln
290 295 300

Ala Trp Asn Asp Ala Pro Cys Asp Lys Thr Phe Leu Phe Ile Cys Lys
305 310 315 320

Arg Pro Tyr Val Pro Ser Glu Pro
325

<210> 3

<211> 10663

<212> DNA

<213> Homo sapiens

<400> 3

tccctgtccc	cactccgcaa	tttgggtgtc	actctacttt	cctccatgac	ttgactgagg	60
gacttgggtg	actcctccct	ggagatgact	cttggcactg	ctggagcatt	tcatacaacc	120
caggtctctg	tccctagtctt	gttgattaaa	cccattggtt	ctctccaagc	ataaggctgg	180
gcccacacat	agtcctcaacc	agagaaaggg	aaaaagtgac	caagctgttt	tcctcctcca	240
tccttatcta	gagaaaaatat	tagaacccca	ggaacacagg	aagaggagaa	actaaaaaaa	300
tgagcacaga	cttcocgttc	tcatacacgc	ggtgattttt	ttggtcacta	ggctcggcca	360
gagttcctct	tctgttttga	agagcagggc	attgtctgag	tgcccgggag	aactgccacc	420
cttctcgact	ggaaaagtgtg	gccaaaggcac	ctgcctgctt	ctctgtcttt	tcctctgcca	480
ggccagggtg	cagatagagc	atccagagggt	ttgcacagga	gaggtgtcta	ggaaagacct	540
gtagatgagc	atgataatag	gaagatggct	cttgactccc	tcctcttcca	aaaagtggga	600
aatgaaactt	cctatgggtt	gggactttta	catccacctc	ctctgaaacc	ccagaaggcc	660
cacagcccac	ggattccatg	ctttttgtct	agcttctcca	tgctggaaac	gcctttcgcc	720
tagcggtagc	tattctacct	acctcagcct	gcatttccat	cacttggagt	aaatgtctcc	780
tgctccagcc	ctctctatgt	ccatggggcca	gccgctaagg	cttctctatc	ctttctcggg	840
accaatgaga	gggcatctgg	ggggctctat	aaagtataac	tcataccaca	aaccagcaga	900
gaaaaaccac	cttctcgacc	gagcggaaat	ggacaagaca	cttcccagcc	acactgtgtg	960
gcggctctct	tcocgcgaat	gcctcaggag	gagacatgga	ccctctgca	gccccttcta	1020
ccaggggtct	gagtaattgt	gattctcttc	tatccccagg	cccagaaata	ggacaggcta	1080
tggtggaagg	gccaggagcc	aggagcagaa	gggcagagat	gattctggag	ttaggctggg	1140
tcaccaagct	gagagtccag	aagtggaaaca	ttctaactct	gtctaagccc	tgatgggaac	1200
aaagctctgc	accacacaga	gccacctgaa	gagagttaga	atctctatgg	acaccacagag	1260
gcaggcgagt	gtgtggagcc	aggccctcat	attcggggact	atcaaatgtt	aaaatttaga	1320
gcagacagca	agctctggaag	ttacagacag	gctgggttagg	gaagcgatgg	gcagagaacaa	1380
atgagacacc	atcttttgcta	cccagtttcc	ttcccagagc	ttcccagctg	ggggtcctct	1440
ctgggtctccc	tcattccagtc	cgagaagcca	ctccagcctg	aggcccggtt	agctggagag	1500
agagcatcaca	gggcctgttaa	ctacaatgag	ctgcacatag	gcattggcgt	agcgtgcagc	1560
cactactctc	ccaagtacca	aaagtgggag	tagccccctga	aagcataggt	agatatcttc	1620
ttggccctgt	taacttaacc	tctaactcctg	gtccctgata	caggctgtca	tcaggctcca	1680
ctggctcagc	ctcagcaggc	ccaggacagc	ttcttttttc	tgccgcocaga	gcttttgttc	1740
attctcgggt	aaggagttaa	tcacacttcc	ccctatgtct	ttgcattgaa	gatgaatgag	1800
aactcttgga	gcactgtgaa	aaaagcagca	gctcttagaa	ggcagatccc	agggcccgca	1860
ctgtgtaaat	gctcgtatgg	catgatcagc	tatcaagctc	ccaagtcact	taccattcac	1920

09787192.031501

acatccatcc	atgcatectc	tatccatcca	ttcatecatt	cacccatcct	ccaattatct	1980
ggaccaaatt	ccttcttttt	ttcttctctt	cctccctctt	tcctctctct	tttccatcat	2040
ttccagaaat	atttgtctag	agtctgcatt	gtgcacaaat	ttctcagatt	gtgcagacttc	2100
ttccaaagcat	gagaaagcgc	ccaggagaaa	atttagtcct	tcattgagcc	caaatcagct	2160
ctctctgccc	ctgagctctt	tgacacatg	gaactctcta	tttaggacta	caagaagga	2220
gagaggagggt	gcggacagag	agagaggaga	gagaaacaga	acgattccta	gtctgaggtt	2280
gctgagcgcc	cctaactttt	ttgaaaaatc	cacatgggtg	aggaagcagg	gagggaacc	2340
caagcttagt	cttatctctt	tgaattccca	agttgcata	ttataaggg	tattgtatca	2400
gcacaaagtc	atataataca	ggaaagcagga	tattctattg	caatgggttg	ctatttggtg	2460
tctctgcttc	caggctcttc	cccaccagtc	tgctctctac	caggatcaga	gtgatgtttc	2520
tacagtgtcc	ctgtccttgt	ctctctctgt	cctgaagtc	tttaattggc	ccatgtcacc	2580
cacaagccag	aggctccagct	ccttgacatg	atcagaagga	ttctcatcat	cagaccccaag	2640
gtgocctctc	aaacccgtgt	gcataccctt	cctgttgga	tgatatattt	acattccagc	2700
aatattgcaa	ctatttaca	tttgccaagc	actccatgt	atcttatgtc	gtgcactttg	2760
gcataatgct	atctttctac	tggaattttt	tttttccatt	tactccattt	catatgcttc	2820
tttgaactgt	gcataattgt	ctctgtgtct	catcacctgt	caattttatt	gtctatgtct	2880
gtctttctca	ccagactatg	agctgtcttg	tgcaaggact	gcgagtattt	catcactgtg	2940
gcctatgcc	tggtagagca	tcagtaccta	gaaggcactc	agcctgtatt	tgtgggggtg	3000
atggatgggt	ggatggagta	cgagagtctt	acaagagaaa	tgggatagg	ttgggacaag	3060
atggttaagt	tatccatgta	acagaccccc	agagaagaca	acaaatggcc	tcttctctgaa	3120
agctcagcat	ctctgaggatg	ggagttaagcc	agacaaggta	tctagtccag	aataagggaag	3180
ttgggatgat	atggtgacct	gctgtgggac	tgacttctgt	tttctctag	ataagagccc	3240
ttggagagcat	aggcagccag	aagcacctgt	gctcccagga	taaggggtgag	cactcaggat	3300
gactgtgcagg	aaggaggccc	ctgatgcgca	cttcaactgt	gacaaacaga	acatctccct	3360
ctggccccga	gcaagccaca	tcgctgtcta	gaacctgtc	cggtgtctgc	gtgcacactt	3420
gccttttgtc	gctccttcaa	cacacatttt	cttctctctc	caacagagcc	tcctcccaag	3480
tccggtccat	ctctggcttc	ggggaaaaca	ccccagctcc	gtgctgcatt	aatctgcctg	3540
acgctgggtc	tggtgcgccc	cgctctgtct	caggccgtcc	tttgaagtc	ctcatgtttc	3600
atgctctggg	cttagctccct	ctctgtgcga	gcgggtctcc	ttcagatgac	gaccactctc	3660
ctctctctcc	ttttttctct	cctgtggctt	tttcaattgt	ctctctctc	ctcttcccat	3720
gtcagtaaac	aggctgtgct	gcccccagta	cggtgagctg	atgctcttct	cctcccaatt	3780
tctggagat	tattgggatt	agcatttgca	catgtgtgct	caaggataca	gttctgtccc	3840
tcaggaaaac	ttgatctagt	tttcatctct	gtgctatttt	ctccccgtcc	acccccacc	3900
aatctgactc	ctgctcctgt	gatctagagc	cttgtggaca	gtcgaatgtc	gtcgaatgtc	3960
tcctgaccct	gtgaaggcgc	gagcctcatg	tatcattggc	cccagctcac	cagatggaga	4020
gcctggccaa	cgagcccaac	gatctccatg	actcagctcc	ctctcccag	gactcgttgg	4080
tgtgtctctc	attctctctg	ccctgtgaag	atcacatcta	gggaggcttc	ctgctcaatt	4140
gatgttgcat	ggtttaactt	ttctctcttt	ctggctctgt	caggcttcat	ggccttttgt	4200
gctgggcag	ccttctctct	ctcgccagg	ctccaggaag	ccagcccaag	ggaccocaag	4260
agctgttggg	tttttttttc	tcctctgtct	gcctcgggac	atattttggc	ttagacttgt	4320
aggctctgag	cagagctccc	cttgccccat	ctagaccccc	tttcttgcta	catcgacttt	4380
tcctcaggcg	tcgctcttga	gocctcttgt	ttatctttga	tcctctctct	ggtaaccagg	4440
cctggagact	gcagcagaat	tgaagtggtg	tggggcaagg	gaaggggaga	aacagttttg	4500
taattctctc	ttccatgttc	tctggagaag	ccacttccag	attagtgctg	gggtctctcc	4560
atggtcacag	aggggcccac	ggacagatgt	gggggagtgg	ttgtctccta	cgagatggcc	4620
actgcagggg	ttcttgaaaa	cagagggatg	gcaaccaagg	gggtgggtct	agggggaact	4680
aagtgtctgc	ggagctgggt	gggtctatgg	agggcacctc	ttatcaaat	ttcctgtaac	4740
actcagaaaa	ttcaggaatg	gtttctaact	ctgtctctg	cttgctctg	aaatcttttc	4800
acagagagcc	tgtgttttat	caactctccc	attatctgta	tcacactgtg	ttcctggcac	4860
atggtagggt	cccatgtcac	gtttgttgta	cgtaaatgaa	ttagtgagg	gttgggggtg	4920
cccattggac	tgtctcttgt	ctttgggaa	cttcagctca	ttcctctccc	tccttttgat	4980
aacctgacaa	caccccccat	cctgtccctg	ggaactcccc	cagctgacac	ctgactcttc	5040
tcaattccag	atcccccggt	tatgggcacc	atatcagatg	taaagaccaa	tgctcagttg	5100
ctgaagagtc	gtgtggacaa	catcagcacc	ctggattctg	aaattaaaaa	gaatagtgc	5160
ggcatggagg	cagctggcgt	tcagatccag	atggtgaaat	agagcctggg	ttatgtgcgt	5220
tctcagcttc	tgaagttaaa	aaccagtgtg	gagaaggcca	acgcacagat	ctcagatctta	5280
acaagaagtt	gggaagaagt	cagtacctta	aatgcccaca	tcccagagtt	aaaaagtgat	5340
ttgggaaaa	ccagtgcttt	aaatacaaa	atccgggacc	tcaggagcag	cttggaaga	5400
atgagcaagt	tgctcaaacg	acaaagttaa	tgaactcaga	aatgagctc	aagctgacca	5460
gtggcccatg	ggatcttacc	tgctccagac	ctgaggccat	tgggctgggt	gggtggggag	5520
gagagtgggg	gcaaaagagg	ggcagccatg	ggctaggaa	ttaaaggag	agggtctgag	5580

gttgggggagg	acttagggggc	tgttaggaga	aaagagacca	gggtccagct	agagctccca	5640
cacaaaagtgt	cagaattgtaa	aagcattagg	ggatgtccac	cctggcccac	acctagtcac	5700
ttcccatcaa	gttcctttct	agagtcagga	ggctcagcca	cttgatcgat	cgatagggag	5760
gttgcttctct	catcatggggg	aagacacctct	ttgtccaacc	tcttgcatca	taacctctcc	5820
agtcaccagag	actctattagg	tctctgtctg	actttcagga	tttgaagagg	tgacctataa	5880
ctcctatacaa	aggaccccaag	gacaccagcg	cacagctcca	tttctgctgt	tctctgagac	5940
ctcattcaag	tgccccccacc	aagccagcat	ccaagaatat	caagaatacc	agcgttcaact	6000
tttactctctt	gttctctaga	tgatattcta	cagggtggtt	ctcaaggctg	gaagtacttc	6060
aagggggaact	tcttactactt	ttctctcatt	ccaagacact	ggatagtgc	cgagcagttc	6120
tgtgtgtcga	ggaattccaca	cctgacctcg	agagtgcaga	ggtagctgct	ggtagctgct	6180
gtgcctatgg	gctctgtgaa	gggggcgtat	gagcactggg	ccaggaggga	tgggcaagat	6240
tatactcgct	gaacaaaaat	ccccaaatat	tgatgacctc	atgaagaagg	attggttctc	6300
agtagcatgt	cgaataaagg	ccggcaaggg	ggctgtgctc	actgtggcta	ctcaaggacc	6360
caggcccaag	gagcttctcat	cttaatatgt	ctccacaatt	gctggggcag	gaaaggggaa	6420
cttgacatat	tgtgcaaaag	ttctgcccg	atttaacata	cctcattttt	ggttacattt	6480
cactagctta	agttatgtta	attaaagtta	gtaagttatg	cagtaactct	ggatggggag	6540
gatcctggat	tccagtaact	ctggaaacagg	gcagagagga	tctactttcc	atgtgcccaag	6600
aaaaaaagaa	atatattgtga	acagccttaa	tgattccaca	tgactcaaga	agtcctctcg	6660
ctggtgaggc	agaaatttggg	caggcccttt	tcatctggga	ggtagggatg	cagagcaggt	6720
cagagcctgg	gctctggccgt	agttctagag	cctgaacctt	gccatctaac	tagtctcagc	6780
agcttgggtg	gaatacccaga	cttcaactgg	caaaccttca	tggccctcac	ttgatgaaac	6840
atgcattgtat	atggcacctg	cctcagagga	aaggagagaa	tgcatatgga	cgactcagca	6900
caattccgta	tgttgattca	gggctcattt	aattcaggta	tctcatatgt	aaccattctc	6960
ttggcgctcg	tgtctggagg	ttcagctgag	gccttctctg	gcttcagcac	ctgcttctcg	7020
agtgccagaa	aggctctgagt	cctgagcttg	ttagctcgag	agcagggaca	ctcataaact	7080
tgggaagatga	aatctgggct	ctgggcaagg	gcaggagaaa	gcttgagagg	ccagtttctg	7140
cagcgcacat	tgtgggtcagg	gctgtcactg	agccaggtgt	aagaacaccc	agagcagtag	7200
gatcaagctc	caagtgtggc	cgcaactctg	cttatctctg	ctttcctaca	ggagtttctg	7260
tataaaaacag	cggggggaact	catctactgg	attggcctga	ctaaagctgg	gatggaaagg	7320
gactgtgtcct	gggtggagtga	cacgccaatt	aaacaaggtc	aaagtgtgag	gtagagccct	7380
ggagccctcc	tgtgccagctc	gactttcccc	ggccatggcc	agggcagtga	gggagtgagg	7440
cgagatgttc	ccatgagaca	gggtttctga	ttcttccctg	tcttagagtg	acaggaaact	7500
tctaacaacca	actcagacaca	accctgtcac	caactggctg	tggacctgag	ccctccacgc	7560
cctctggggt	tggcacaaca	ggccttctac	ctggccagct	tcagggatct	gtctcatagt	7620
ctaggttcttc	acagttgtggg	tttgtgtagg	gacttgaagg	tgggtgtctg	gtttgtccctg	7680
gacttggggc	atgtgaaaag	ttagaggtcg	aagttctcaac	agtcctcttc	ctctgaggct	7740
tgggtgcaga	catttgcctat	gccattccct	aggacaaaag	cttgggttga	gttaactcat	7800
ttcttcaact	gaaataagbt	ctttttgatt	ttccactttg	taaatccact	tttttccccc	7860
ctctgttagtg	gttctgtgatt	ccagggtgagc	ccaacaatgc	tgggaacaat	gtaactctgt	7920
gcaatataaa	ggctcctcaca	cttcaggcct	ggaatgatgc	cccatgtgac	aaaaagtttc	7980
tttctatttg	taagcgaccct	tatgtcccat	cagaaaccgtg	acaggacagg	ctcccaagct	8040
caactcttga	gctccaacgc	ttgttaaaca	taggagaaatg	cctcttctct	cccagactgc	8100
caggatgact	tgtccagctta	atttttcttg	cttcaaaaat	gtcccacagt	ggcattctgg	8160
agtcctgtct	tcttggctgtg	aaattctctg	acgtcttgga	ggcagctgga	atggaaaagg	8220
gaattcagtg	taaaagtgga	ggggtgggta	gagaggattt	agaagttcca	atgtccctgc	8280
taaggaggat	caagaccctg	aatccggcat	aacacccctg	ggttttccac	tctttcagag	8340
aaacctcagc	tcttccatcat	caaatgaact	ccagagcaac	caagcaactc	ctctgatatt	8400
gtcatccagg	gttctttctg	gccaaacccc	ctagaatttc	catgttctg	cttagctgtg	8460
ctggcagctca	gcagctggct	gtgtttgcag	tgcaaatgac	tctgttctg	gaaactctgc	8520
tcatgttatg	ttcccaagtgt	tttctctc	cacatcatct	aaagcctgaa	ccggttcttc	8580
tttgtgtcaa	gtcagttggct	gacacggact	tgtatctcct	tcagagctgt	gctggcaccc	8640
agcctccctt	ctccttccac	ttccttagta	cactggagtg	ccagagccctg	ctctccaccc	8700
agcgttccat	cagccctctg	cctcactctc	ccggcacctc	ctcctccttc	tgcatttctc	8760
atcttctgtg	gtcttttgca	tgggaagcag	cttctagatc	cttctgaatc	tacacctcca	8820
gcttctcagc	aaataaagtgt	tgcttgggtc	aaggactcac	tccaagtcca	cttttttact	8880
tctgtgtgtc	caggtgaaata	tgtgggaaaag	cgagttctct	tgtgtggaca	tggagtctta	8940
gggtatcttc	aggaacaactc	tggggagtca	aaaataacaa	ggactgggga	agttccagtc	9000
ctggaaaatgc	cacaaaattgt	gaccagtact	tattcttagt	ttttatataa	gtagagcaag	9060
gtctccaatg	tcaogtatgt	ggtagatctt	ctcttctgtt	actgcacaat	ctcttagtgt	9120
atagctcaat	ccccagaacac	aagttctcagc	aggttcccca	ctcttcacag	agaccagagt	9180
ccacaggcat	tacttccaaa	tcccaagtc	agtggctgaa	agggcaatcc	agggcagcgc	9240

caccacagag aggagaggag ggtggagtga gcacagggtc tcattaagggt cctcagga 9300
 agatgcttcc ttaataaact gtaaccagca gtgtgtgtgt ctgggtgcaa atgggtcaca 9360
 gctgagggca cagggttgta ttgtaagacc tgaaatacca cgtgctgctg tgacatttta 9420
 tgctctacag ggccccaaag acctaacctc gagttccctg cctctcaccg gatatactct 9480
 tgccctcggt cccacactgg ctaatttctc atcatctgga ccagctgcgt gccaccaggt 9540
 ttctacttta atgggtttca cttctctgcc agcctgagaa actattcaaa caagccaatc 9600
 acatctctct acaggaatcc ggggcactct atccttttat tactacaagg cctgctctcc 9660
 acagccctgg ctggttcaat ctgctcctga gggtagcccc atgtggccct gtgtggctta 9720
 tgatatcttt cccacagtag ctgtatttgt gactagtaaa ctgctgccag tctccactgc 9780
 acagtgtcaa atgtcttgtt ttggccatct tgtctattt agagcagggg atccctccct 9840
 caccaatgga ggaataggga ggtgacaaga acaaggctgg tcggggatct ctggttggtt 9900
 ttggcaaga gatgagctgg gaaatcaga ccatttctct gggaaagaat ttgaaccagg 9960
 aaatagcaag aggatgaggg tgtaaacaaa aggaagtga gctggaaggc actgagttaa 10020
 gagaaaggct ggagggggcc tcacgtggca ttggaagaaa ctgacaatga gcagaagcta 10080
 tgaggcaggg gaaagacatg aatacaggcc tggcatggtg gctcatgcct ataatcccag 10140
 cactttggga ggctgaagtg ggtgggtcac ctgaggtcac gagtctgaga cagcctggcc 10200
 aacatgggta aacccatct ctactaaaaa tacaaaaatt agccgggcat ggtggtgggc 10260
 acctgtaatt ccaactactt gggagactga ggcaggagaa ttgcttgaat ctgggaggca 10320
 gaggttgtag tgagctgaga tcccaccact gcactccagc ctgggcaaca gggcaagact 10380
 ttgtttcaaa aaaaaagaag tgactgcaga ggattatagt tggcagagaa aagagaacgg 10440
 ctacaggag tcgcaatgga ggtcccggag ggcagcctga agggctccgg ctgctccgt 10500
 tcccagggtc gctcagatc ctcccagccc ttctgatctc cctggtttct gtgcatgggg 10560
 accttcagag gctgtgctcc tgacccaac cattgtttt tcttgaaact gaaagagcct 10620
 gagtcatgta ggatgtgttt ttatctggag tctgtgcccc agc 10663

<210> 4
 <211> 120
 <212> DNA
 <213> Homo sapiens

<400> 4
 ggcagccaga agcacctgtg ctcccaggat aagggtagac actcaggatg actgtggaga 60
 aggaggcccc tgatgcgcac ttactgttg acaaacagaa catctccctc tggccccagg 120

<210> 5
 <211> 117
 <212> DNA
 <213> Homo sapiens

<400> 5
 agctcctccc caagtccggt ccactctctg tccccgggaa aacaccaca gtcctgtctg 60
 cattaatctg cctgacgctg gtctgtgctg cctcctgctc gctgcaggcc gtccttt 117

<210> 6
 <211> 375
 <212> DNA
 <213> Homo sapiens

<400> 6
 atccccgggt tatgggcacc atatcagatg taaagaccaa tgtccagttg ctgaaaggto 60
 gtgtggacaa catcagcacc ctggattctg aaattaaaaa gaatagtgc gcatggagg 120
 cagctggcgt tcagatccag atggtgaatg agagcctggg ttatgtgcgt tctcagttcc 180
 tgaagttaaa aaccagtgtg gagaaggcca acgcacagat ccagattctta acaagaagt 240
 gggagaagt cagtacctta aatgccccaa tcccagagtt aaaaagtgat ttggagaag 300
 ccagtgcttt aaatacaaa atccgggcaac tccagggcag cttggagaat atgagcaagt 360
 tgctcaaacg acaaa

<210> 7
 <211> 152
 <212> DNA
 <213> homo sapiens

<400> 7
 atgatattct acaggtgggt tctcaaggct ggaagtactt caaggggaac ttctattact 60
 ttctctcat tccaagacc tggatatagt ccgagcagtt ctgtgtgtcc aggaattcac 120
 acctgacctc ggtgacctca gagagtgage ag 152

<210> 8
 <211> 119
 <212> DNA
 <213> Homo sapiens

<400> 8
 gagtttctgt ataaacacgc ggggggactc atctactgga ttggcctgac taaagcaggg 60
 atggaagggg actgggtcctg ggtggatgac acgccattca acaaggtcca aagtgcgag 119

<210> 9
 <211> 985
 <212> DNA
 <213> Homo sapiens

<400> 9
 gttctgggatt ccaggtgagc ccaacaatgc tgggaacaat gaacactgtg gcaatataaa 60
 ggctccctca cttcaggcct ggaatgatgc cccatgtgac aaaacgtttc ttttcatttg 120
 taagcgaccc tatgtcccat cagaaccgtg acaggacagg ctcccgaagt cactctttga 180
 gctccaacgc ttgttaaaaca tgaggaaatg cctctttctt cccagactc caggatgact 240
 ttgcacgtta atttttcttg ctcaaaaatt gtcccacagt ggcattctgg agtcgctctg 300
 tcttggtctg aaattctctg acgtcttgga ggcagctgga atggaaagga gaattcaggt 360
 taaagtggga ggggtgggta gagaggattt agaagttcca attgccctgc taaggaggat 420
 caagaccctg aatccggcac aacaccctgg ggttttccac tcttccagag aaacctcagc 480
 ttcatcacat caaagtact ccagagcaac caagcaattc tctgatatt gtcatccagg 540
 gcttttcttg gccaaaacccc ctagaatttc catgtctctg cttagctgtg ctggcagcta 600
 gcagctggct gtgtttgcag tgcataatgc tctgttcttg gaaatcctgc tcatgggtatg 660
 tcccagctgg ttctcttcac cacatcatct aaagcctgaa cccgttcttc tctggttcaa 720
 gtcagtggtc gacacggact tgtatctcct tcagagctcg gctggcaccc agcctccctt 780
 ctctctccac tcccttagta cactggagtg ccgagccctg ccttccaccc agcgtccatc 840
 cagccctgt cctcactctc ccggcacctc ctctcctctc tgcatttctc atcttctgtg 900
 gtctgtgca tgggaagcag ccttcagtgc ctctcatgaa tcaccttcca gcttccctcag 960
 aataaaatgc tgcctgggtc aagga 985

<210> 10
 <211> 1756
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (266)..(1243)

<400> 10
 ggctcttttg ctgcttttgt gtcttcagtt aagatgatta aaatgtctgt gtaccagga 60
 agtagtcagg tggcttcctc atcaaaagccc aattcttagt caaaaggatg aagctgtggc 120
 ctctgccttg ctaatcgtct aagccaagaa ctgggaacct gggcatgaca aagtggcctg 180

ctttttggct	catttccgt	tctttctcgg	atgaaaaggt	ccttggggag	acggatatcc	240						
agctttttcct	aagccagagg	cagag	atg	ttg	gag	gag	gct	ccc	gaa	gcg	cac	292
		Met	Leu	Glu	Glu	Ala	Pro					
		1					5					
ttc	aca	gtg	gac	aaa	cag	aac	atc	tct	ctc	tgg	cct	340
Phe	Thr	Val	Asp	Lys	Gln	Asn	Ile	Ser	Leu	Trp	Pro	
10				15						20		
ccc	aag	caa	gat	ctg	tct	cca	gtt	ctg	agg	aaa	cct	380
Pro	Lys	Gln	Asp	Leu	Ser	Pro	Val	Leu	Arg	Lys	Pro	
				30					35			
gtg	gcc	ttc	acc	tgc	ctg	gca	ttg	gtg	ctg	gtc	acc	436
Val	Ala	Phe	Thr	Cys	Leu	Ala	Leu	Val	Leu	Val	Thr	
			45					50				
cag	gct	gtt	ttc	tat	cct	agg	ttg	atg	ggc	aaa	ata	484
Gln	Ala	Val	Phe	Tyr	Pro	Arg	Leu	Met	Gly	Lys	Ile	
		60					65				70	
agt	gat	gcc	cag	atg	ttg	aaa	ggg	cgt	gtg	gac	aac	532
Ser	Asp	Ala	Gln	Met	Leu	Lys	Gly	Arg	Val	Asp	Asn	
	75					80					85	
ggg	tct	gat	ctt	aag	act	gaa	aga	ggg	cgt	gtg	gac	580
Gly	Ser	Asp	Leu	Lys	Thr	Glu	Arg	Gly	Arg	Val	Asp	
90				95						100		
cag	atg	cag	ata	gtg	aac	acc	acc	ctc	aag	agg	gtg	628
Gln	Met	Gln	Ile	Val	Asn	Thr	Thr	Leu	Lys	Arg	Val	
			110						115			
ctg	tct	ttg	gaa	acc	agc	atg	aag	ata	gcc	aat	gat	676
Leu	Ser	Leu	Glu	Thr	Ser	Met	Lys	Ile	Ala	Asn	Asp	
			125					130				
tta	aca	atg	agc	tgg	gga	gag	gtt	gac	agt	ctc	agt	724
Leu	Thr	Met	Ser	Trp	Gly	Glu	Val	Asp	Ser	Leu	Ser	
		140				145					150	
gaa	ctg	aaa	aga	gat	ctg	gat	aaa	gcc	agc	gcc	ttg	772
Glu	Leu	Lys	Arg	Asp	Leu	Asp	Lys	Ala	Ser	Ala	Leu	
	155					160					165	
caa	gga	cta	cag	aac	agc	ttg	gag	aat	gtc	aac	aag	820
Gln	Gly	Leu	Gln	Asn	Ser	Leu	Glu	Asn	Val	Asn	Lys	
170				175					180			
cag	agt	gac	att	ctg	gag	atg	gtg	gct	cga	ggc	tgg	868
Gln	Ser	Asp	Ile	Leu	Glu	Met	Val	Ala	Arg	Gly	Trp	
			190						195			
ggg	aac	ttc	tat	tac	ttt	tca	cgc	acc	cca	aag	acc	916
Gly	Asn	Phe	Tyr	Tyr	Phe	Ser	Arg	Thr	Pro	Lys	Thr	
		205						210			215	
gag	cag	ttc	tgt	att	tct	aga	aaa	gct	acc	ctg	acc	964
Gly	Gln	Phe	Cys	Ile	Ser	Arg	Lys	Ala	His	Leu	Thr	
										Ser	Val	

220 225 230

gaa tcg gaa caa aag ttt ctc tac aag gca gca gat gga att cca cac 1012
 Glu Ser Glu Gln Lys Phe Leu Tyr Lys Ala Ala Asp Gly Ile Pro His
 235 240 245

tgg att gga ctt acc aaa gca ggg agc gaa ggg gac tgg tac tgg gtg 1060
 Trp Ile Gly Leu Thr Lys Ala Gly Ser Glu Gly Asp Trp Tyr Trp Val
 250 255 260 265

gac cag aca tca ttc aac aag gag caa agt agg agg ttc tgg att cca 1108
 Asp Gln Thr Ser Phe Asn Lys Glu Gln Ser Arg Arg Phe Trp Ile Pro
 270 275 280

ggg gaa ccc aac aac gca ggg aac aac gag cac tgt gcc aat atc agg 1156
 Gly Glu Pro Asn Asn Ala Gly Asn Asn Glu His Cys Ala Asn Ile Arg
 285 290 295

gtg tct gcc ctg aag tgc tgg aac gat ggt ccc tgt gac aat aca ttt 1204
 Val Ser Ala Leu Lys Cys Trp Asn Asp Gly Pro Cys Asp Asn Thr Phe
 300 305 310

ctt ttc atc tgc aag agg ccc tac gtc caa aca act gaa tgacagatct 1253
 Leu Phe Ile Cys Lys Arg Pro Tyr Val Gln Thr Thr Glu
 315 320 325

ggcctgagct cggcatctgt ggggcaacag tgacctggct gaagagatgt ctctctccct 1313

gagggtccaa gattgctctg tactttacgtt tttttcttgc ttgaaaattg tcccaaacac 1373

agcctgtggt ctttctgtct tggtctggcag ttctctgctc ctggaggcct tggaggagct 1433

tgggttaaac ggggtaggac ctgaaaaggg tgtagcagtc cttactgccc aggcgaggca 1493

ggtcagcaca ccaaacaggt tgtttagatt ttcttgatcc ttctcagaag ccttggtctga 1553

ccatataaaa gctacattca aatatgacca gtatttgagg agacagacat gcccaaat 1613

aaccatgata caatttatca aacatgtatt agaacacctc atggtatggt caaaatagta 1673

aatatgttgt ttttatgtgc ctattgcaaa taaatgtaaa gacttaaaaa aaaaaaaaaa 1733

aaaaaaaaaa aaaaaaaaaa aaa 1756

<210> 11
 <211> 326
 <212> PRT
 <213> Mus musculus

<400> 11
 Met Leu Glu Glu Ala Pro Glu Ala His Phe Thr Val Asp Lys Gln Asn
 1 5 10 15

Ile Ser Leu Trp Pro Arg Glu Pro Pro Pro Lys Gln Asp Leu Ser Pro
 20 25 30

Val Leu Arg Lys Pro Leu Cys Ile Cys Val Ala Phe Thr Cys Leu Ala
 35 40 45

Leu Val Leu Val Thr Ser Ile Val Leu Gln Ala Val Phe Tyr Pro Arg

09787192-031501

50	55	60
Leu Met Gly Lys Ile Leu Asp Val Lys Ser Asp Ala Gln Met Leu Lys 65 70 75 80		
Gly Arg Val Asp Asn Ile Ser Thr Leu Gly Ser Asp Leu Lys Thr Glu 85 90 95		
Arg Gly Arg Val Asp Asp Ala Glu Val Gln Met Gln Ile Val Asn Thr 100 105 110		
Thr Leu Lys Arg Val Arg Ser Gln Ile Leu Ser Leu Glu Thr Ser Met 115 120 125		
Lys Ile Ala Asn Asp Gln Leu Leu Ile Leu Thr Met Ser Trp Gly Glu 130 135 140		
Val Asp Ser Leu Ser Ala Lys Ile Pro Glu Leu Lys Arg Asp Leu Asp 145 150 155 160		
Lys Ala Ser Ala Leu Asn Thr Lys Val Gln Gly Leu Gln Asn Ser Leu 165 170 175		
Glu Asn Val Asn Lys Leu Leu Lys Gln Gln Ser Asp Ile Leu Glu Met 180 185 190		
Val Ala Arg Gly Trp Lys Tyr Phe Ser Gly Asn Phe Tyr Tyr Phe Ser 195 200 205		
Arg Thr Pro Lys Thr Trp Tyr Ser Ala Glu Gln Phe Cys Ile Ser Arg 210 215 220		
Lys Ala His Leu Thr Ser Val Ser Ser Glu Ser Glu Gln Lys Phe Leu 225 230 235 240		
Tyr Lys Ala Ala Asp Gly Ile Pro His Trp Ile Gly Leu Thr Lys Ala 245 250 255		
Gly Ser Glu Gly Asp Trp Tyr Trp Val Asp Gln Thr Ser Phe Asn Lys 260 265 270		
Glu Gln Ser Arg Arg Phe Trp Ile Pro Gly Glu Pro Asn Asn Ala Gly 275 280 285		
Asn Asn Glu His Cys Ala Asn Ile Arg Val Ser Ala Leu Lys Cys Trp 290 295 300		
Asn Asp Gly Pro Cys Asp Asn Thr Phe Leu Phe Ile Cys Lys Arg Pro 305 310 315 320		
Tyr Val Gln Thr Thr Glu 325		

09787192-034501

DECLARATION AND POWER OF
ATTORNEY FOR PATENT APPLICATION

Attorney's Docket No. **SF0820K2**

As a below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name;

I believe I am the original, first sole inventor (if only one name is listed below) or an original,
first

and joint inventor (if plural names are listed below) of the subject matter which is claimed and
for which a patent is sought on the invention entitled:

"ANTIBODIES TO MAMMALIAN DC ANTIGEN AND USES"

the specification of which

☒ is attached hereto.

☐ was filed on _____ as Application Serial No. _____

and was amended on _____ (if applicable).

☐ was filed on September 23, 1999 as PCT International Application
No. PCT/US99/22269

I hereby state that I have reviewed and understand the contents of the above-identified
specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the patentability of this
application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, §119(a)-(d) of any
foreign application(s) for patent or inventor's certificate listed below and have also identified
below any foreign application for patent or inventor's certificate having a filing date before that
of the application on which priority is claimed:

Prior Foreign Application(s):		Priority Claimed
<u>98 402 374.7</u>	<u>Europe</u>	<u>25 September 1998</u>
(Number)	(Country)	(Day/Month/Year Filed)
		Yes or No

Prior Foreign Application(s):		Priority Claimed
<u>99 400 394.5</u>	<u>Europe</u>	<u>18 February 1999</u>
(Number)	(Country)	(Day/Month/Year Filed)
		Yes or No

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States
provisional application(s) listed below:

(Application Number)	(Filing Date)
----------------------	---------------

I hereby claim the benefit under Title 35, United States Code, §120 of any United States
application(s) listed below and, insofar as the subject matter of each of the claims of this
application is not disclosed in the prior United States application in the manner provided by the
first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose
material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which
occurred between the filing date of the prior application and the national or PCT international
filing date of this application:

(Application Serial No.) (Filing Date) (Status - patented, pending, abandoned)

Power of Attorney: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. (List name and registration number.)

Carl W. Battle	Reg. No. 30731	Susan Lee	Reg. No. 30653
Edwin P. Ching	Reg. No. 34090	Anita W. Magatti	Reg. No. 29825
Eric S. Dicker	Reg. No. 31699	Joseph T. Majka	Reg. No. 30570
Cynthia L. Foulke	Reg. No. 32364	Arthur Mann	Reg. No. 35598
Robert A. Franks	Reg. No. 28605	Edward H. Mazer	Reg. No. 27573
Kenneth M. Goldman	Reg. No. 34174	Jaye P. McLaughlin	Reg. No. 41211
James M. Gould	Reg. No. 33702	Richard B. Murphy	Reg. No. 35296
Richard J. Grochala	Reg. No. 31518	James R. Nelson	Reg. No. 27929
Henry S. Hadad	Reg. No. 35888	Immac J. Thampoe	Reg. No. 36322
Thomas D. Hoffman	Reg. No. 28221	Paul A. Thompson	Reg. No. 35385
Henry C. Jeanette	Reg. No. 30856	Joanne P. Will	Reg. No. 35737
Palaiyur S. Kalyanaraman	Reg. No. 34634	Donald W. Wyatt	Reg. No. 40879
Gerald P. Keleher	Reg. No. 43707		

Send Correspondence to: Cynthia L. Foulke <u>Schering-Plough Corporation</u> <u>Patent Department, K-6-1, 1990</u> <u>2000 Galloping Hill Road</u> <u>Kenilworth, New Jersey 07033-0530</u>	Direct Telephone Calls to: Name: Cynthia L. Foulke Telephone No.: (908) 298-2135 Facsimile No.: (908) 298-5388
--	---

FULL NAME OF 1ST OR SOLE INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
	Duvert-Frances	Valerie	
RESIDENCE & CITIZENSHIP	CITY Tasin La Demi-Lune	STATE OR FOREIGN COUNTRY France	COUNTRY OF CITIZENSHIP France
POST OFFICE ADDRESS	POST OFFICE ADDRESS 1-3 chemin de la Pomme Résidence les Hespérides - Allée E	CITY 69160 Tasin La Demi-Lune	STATE & ZIP CODE/COUNTRY France

FULL NAME OF 2ND JOINT INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
	Pin	Jean-Jacques	
RESIDENCE & CITIZENSHIP	CITY Saint Bonnet de Mure	STATE OR FOREIGN COUNTRY France	COUNTRY OF CITIZENSHIP France
POST OFFICE ADDRESS	POST OFFICE ADDRESS 94 RN6 69720 Saint Bonnet de Mure	CITY 69720 Saint Bonnet de Mure	STATE & ZIP CODE/COUNTRY France

FULL NAME OF 3RD JOINT INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME
	Valladeau	Jenny	
RESIDENCE & CITIZENSHIP	CITY Lyon	STATE OR FOREIGN COUNTRY France	COUNTRY OF CITIZENSHIP France
POST OFFICE ADDRESS	POST OFFICE ADDRESS 10 rue Camille Roy	CITY 69007 Lyon	STATE & ZIP CODE/COUNTRY France

09787192-03504

W

2W

3W

40

FULL NAME OF 4TH JOINT INVENTOR	FAMILY NAME Clair	FIRST GIVEN NAME Valerie	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY Lyon	STATE OR FOREIGN COUNTRY France	COUNTRY OF CITIZENSHIP France
POST OFFICE ADDRESS	POST OFFICE ADDRESS 21 Avenue Lacassagne	CITY 69003 Lyon	STATE & ZIP CODE/COUNTRY France







SW

FULL NAME OF 5TH JOINT INVENTOR	FAMILY NAME Saeland	FIRST GIVEN NAME Sem	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY Lyon	STATE OR FOREIGN COUNTRY France	COUNTRY OF CITIZENSHIP Norway
POST OFFICE ADDRESS	POST OFFICE ADDRESS 6 rue Soeur Bouvier	CITY 69005 Lyon	STATE & ZIP CODE/COUNTRY France

SW

FULL NAME OF 6TH JOINT INVENTOR	FAMILY NAME Lebecque	FIRST GIVEN NAME Serge	SECOND GIVEN NAME J.E.
RESIDENCE & CITIZENSHIP	CITY Civrieux d'Azergues	STATE OR FOREIGN COUNTRY France	COUNTRY OF CITIZENSHIP Belgium
POST OFFICE ADDRESS	POST OFFICE ADDRESS 54 Chemin du Marand	CITY 69380 Civrieux d'Azergues	STATE & ZIP CODE/COUNTRY France

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of First Inventor 	Signature of Second Inventor 	Signature of Third Inventor 
Date 10/04/93	Date 10.04.93	Date 10/01/99
Signature of Fourth Inventor 	Signature of Fifth Inventor 	Signature of Sixth Inventor 
Date 10.04.99	Date 10.01.99	Date 10.01.99

CERTIFIÉ SINCÈRE ET VÉRITABLE PAR

M^e 82-1-11-16-51- notaire

associé à ECULLY (Rhône), le

signature de Valérie JUVERT - Sem Saeland SW -

apposée ci-dessus Semy VACHEAU - Valérie CLAIR

ECULLY le 6/10/99 Sem SAELAND - Serge LEBECQUE

